

Due to a submission error, the article, “A Combined Analysis of 2 Randomized Clinical Studies of Tretinoin Gel 0.05% for the Treatment of Acne” (*Cutis*. 2009;83:146-154), reported the incorrect number of participants for the skin-related adverse event of skin burning sensation in the vehicle group (Table 5). Consequently, the entire table is reprinted below in its corrected form. *Cutis*[®] makes every possible effort to ensure accuracy in its articles and apologizes for the mistake.

Table 5.

Combined Incidence of Skin-Related Adverse Events^a

Adverse Event	Treatment Group			P Value ^b
	Tretinoin Gel 0.05% (n=674)	Tretinoin Gel Microsphere 0.1% (n=376)	Vehicle (n=487)	
Skin related, n (%)	208 (31)	196 (52)	25 (5)	<.001
Exfoliative dermatitis	37 (5)	80 (21)	4 (1)	<.001
Dry skin	109 (16)	112 (30)	8 (2)	<.001
Erythema	47 (7)	67 (18)	1 (<1)	<.001
Scaly rash	14 (2)	29 (8)	1 (<1)	<.001
Skin burning sensation	53 (8)	57 (15)	8 (2)	<.001

^aCommon adverse events experienced by $\geq 5\%$ of participants in at least 1 treatment group.

^bOverall P value derived from Fisher exact test.