

Excoriations and ulcers on the arms and legs

Richard P. Usatine, MD

Department of Family and Community Medicine, University of Texas Health Science Center at San Antonio

Miguel A. Saldaña-Arregui, BA

School of Medicine, University of Texas Health Science Center at San Antonio

A 55-year-old woman came to the clinic complaining of severe itching on her arms and legs. Although she itched throughout the day, it became intolerable at night, disrupting her sleep. She would sometimes scratch her arms and legs until exhaustion but could find no relief. Being outside in warm and sunny weather aggravated the problem. She had used moisturizers, emollients, and topical corticosteroids, but they only alleviated the itching temporarily. The itching began 10 months earlier, just after she finalized the divorce from her husband of 20 years.

Examination of the skin revealed numerous excoriations with ulcerations and xerosis on the arms and left leg (**Figure 1 and 2**). The excoriations were located extensively from the dorsum of

her left foot to above the knee and bilaterally from the wrist to above the elbow. They also showed signs of infection. The patient admitted they were self-inflicted. The patient's right leg had been amputated 5 years before after a car accident, and she wore a prosthetic leg. Examination of other areas showed nothing remarkable.

The patient readily admitted to a great deal of psychological distress. She described feeling depressed since her divorce. She has had difficulty securing a full-time job and has high anxiety about being able to pay her rent and bills.

■ WHAT IS THE DIAGNOSIS?

■ WHAT IS THE TREATMENT STRATEGY?

FIGURE 1 Excoriations on the arms...



Multiple excoriations and ulcers cover the patient's left leg and both her arms.

FIGURE 2 ...and the left leg



Detail of the excoriations and ulcers on the left leg. The right leg is a prosthesis.

FIGURE 3 Detail of lesions on the arm

Crusting and redness indicative of localized infection.

■ DIFFERENTIAL DIAGNOSIS: PSYCHODERMATOLOGIC DISORDER

The patient's history and physical examination points to a psychodermatologic disorder. Psychodermatologic disorders are conditions involving an interaction between the mind and skin and are classified as:³

- *Psychophysiological disorders:* skin disorders worsened by emotional stress
- *Primary psychiatric disorders:* usually caused by psychological conditions with self-induced skin damage
- *Secondary psychiatric disorders:* psychological problems developed as a consequence of a disfiguring skin disorder, which negatively effects self-esteem and body image.³

This patient's excoriations and ulcers are due to self-mutilation. The differential diagnosis includes psychogenic parasitosis, factitial dermatitis, and neurotic excoriations. These 3 skin conditions are primary psychiatric symptoms, and proper diagnosis revolves around being able to assess the dermatologic features and associated psychological disorder.²

■ SUBMITTING IMAGES TO PHOTO ROUNDS

Do you have images (slides, prints, digitized photos) of compelling clinical cases of interest to family physicians? We would like to publish them, along with a brief description of the clinical presentation and a diagnostic question for readers. The case should include information on the differential diagnosis and treatment, the latter applying an evidence-based approach supported by current references. Submit electronic files to usatine@uthscsa.edu, or send high-quality slides and prints to:

Submissions: Richard P. Usatine, Editor, Photo Rounds, University of Texas Health Sciences Center at San Antonio, Dept of Family and Community Medicine, MC 7794, 7703 Floyd Curl Drive, San Antonio, TX 78229-3900.

Psychogenic parasitosis

Also known as delusional parasitosis, psychogenic parasitosis is a psychodermatologic disorder in which patients believe they are infested with parasites. Patients with this disorder report seeing or feeling parasites on their skin, and they damage their skin in an attempt to remove them. Patients create excoriations and ulcers on easily reached areas, usually the ears, eyes, nose, and extremities. They often present with what is termed the "matchbox sign," in which they bring containers filled with "small bits of excoriated skin, dried blood, debris, or insect parts as proof of infestation."²

Women over the age of 50 years are more often affected with psychogenic parasitosis, and the disorder is associated with anxiety, depression, and hypochondriasis.

Factitial dermatitis

Factitial dermatitis, also known as dermatitis artefacta, is a psychodermatologic disorder in which patients damage their skin but deny their self-involvement. This disorder encompasses a wide range of lesions, including blisters, cuts, ulcers, and burns. Patients often are unable to describe how the lesions evolved. Lesions exhibit bizarre patterns not characteristic of any disease.

Young adults and adolescents are more commonly affected, and it is 4 times more common among women than men. Psychological disorders involved with factitial dermatitis include personality disorders and posttraumatic stress disorder.²

Neurotic excoriations

Neurotic excoriations, sometimes referred to as neurodermatitis, are a result of a psychodermatologic

disorder in which patients inflict excoriations and ulcers on their skin and admit to their involvement.³ The condition is characterized by excoriations similar in size and shape that are localized on areas easily reached by the patient, such as the arms, legs, and upper back.

The lesions may present in various stages, varying from dug-out ulcers to ulcers covered with crusts and surrounded by erythema to areas receding into depressed scars. These lesions are a result of repetitive scratching and digging by the patient, usually without an underlying physical pathology but sometimes initiated by pruritus.^{1,2}

Studies show the condition primarily affects women, with a mean onset between the ages of 30 to 45 years.¹ Common psychiatric problems associated with neurotic excoriations include significant social stress, depression, anxiety, and obsessive-compulsive disorder.

Diagnosis: Neurotic excoriations due to depression and stress

The patient's history and physical examination suggests a diagnosis of neurotic excoriations due to the characteristics of the excoriations and ulcers on her arms and leg, her admission of their self-inflicted nature, and the associated depression and psychosocial stress.

■ LABORATORY TESTS

Although there are no available laboratory tests to confirm a positive finding of neurotic excoriations, tests could be performed to disprove any systemic causes of pruritus and the resulting dermatological damage.⁵ These tests include complete blood count with differential, chemistry profile, determination of thyroid-stimulating hormone levels, fasting plasma glucose level, and skin biopsies.^{1,4}

Corresponding author: Richard P. Usatine, MD, University of Texas Health Sciences Center at San Antonio, Dept of Family and Community Medicine, MC 7794, 7703 Floyd Curl Drive, San Antonio, TX 78229-3900.

Laboratory tests for systemic causes of pruritus should be based on the patient's physical and history, avoiding a broad approach.

■ TREATMENT: ADDRESS THE SKIN AND THE PSYCHE

The treatment of neurotic excoriations requires a dual approach, addressing both the dermatological problems and the underlying psychological disorder.³ Supportive dermatologic care is necessary to avoid secondary complications and to ensure that the patient feels supported.³

Dermatologic care

Topical steroids can be helpful to decrease the pruritus and inflammation. The steroid strength should be chosen based upon the severity of the lesions and the thickness of the skin. Steroid ointments are preferred to creams when there are skin ulcers and deep excoriations.

If there is significant crusting or exudate, there is probably a secondary bacterial infection and oral antibiotics are indicated (Figure 3). A first-generation cephalosporin or dicloxacillin should provide adequate coverage against the most likely organisms, *Staphylococcus aureus* and *Streptococcus pyogenes*.

Psychological care

The selective serotonin reuptake inhibitors produce a strong antipruritic response in patients with neurotic excoriations. Doxepin is a tricyclic antidepressant with one of the most powerful antihistamines for pruritus. When prescribing doxepin, limit the amount of medication dispensed at one time to minimize the risk of suicide.³ The patient should trim the fingernails to reduce the amount of damage caused by scratching and digging.

Treating the underlying psychological disorder requires supportive and empathic counseling. It may be necessary to collaborate with other mental health specialists. Management options include psychotropic medication, stress management courses, and referral to a psychiatrist.

New Options in Hormone Therapy: Safety, Efficacy, and What Your Patients Need to Know

TOPICS

■ Cardiovascular Disease and Hormone Therapy: What the data show

David F. Archer, MD

*Professor of Obstetrics and Gynecology
Eastern Virginia Medical School*

■ Assessing Risks and Benefits of Hormone Therapy for the Individual Patient:

**Breast cancer, osteoporosis,
and cognitive decline**

James A. Simon, MD

*Clinical Professor of Obstetrics and Gynecology
George Washington University School of Medicine*

■ New Hormone-Therapy Formulations and Routes of Delivery:

Meeting the needs of your patients in the post-WHI world

Vivian Lewis, MD

*Strong Fertility and Reproductive Science Center
University of Rochester Medical Center*

**Look for the supplement
at www.jfponline.com.**



The American Society for Reproductive Medicine is accredited by the ACCME to provide continuing medical education for physicians.



Supported by an unrestricted educational grant from Solvay Pharmaceuticals, Inc.



Medical Education Resources, a nonprofit medical education company, is accredited by the Accreditation Council for Pharmacy Education as a provider of CEU.

This supplement is based on symposium presentations from ASRM's 2003 Annual Meeting in San Antonio, Texas.

Patients with psychodermatologic disorders frequently resist referral to mental health professionals.³ If this is the case, family physicians are well positioned to help patients with psychological problems.

Alternative courses of treatment include hypnosis to disrupt the itch-scratch cycle, and acupuncture and supportive therapy to reduce underlying stressors.¹

■ PATIENT'S OUTCOME

By showing support and concern for the patient's health during the visit, the physician strengthened the relationship with the patient, who felt comfortable disclosing many of her concerns and troubles. She received prescriptions for cephalexin 500 mg orally 3 times daily, 0.1% triamcinolone ointment to be applied twice daily, and doxepin 25 mg once nightly. She also received a referral for counseling.

At first she refused to trim her fingernails, but as she began to see that her ulcers were self-inflicted, she reconsidered and agreed. In fact, she asked to borrow our nail clippers before leaving the office.

REFERENCES

1. Cyr PR, Dreher GK. Neurotic excoriations. *Am Fam Physician* 2001; 64:1981-1984.
2. Habif TP. *Clinical Dermatology*. 4th ed. New York: Mosby; 2004.
3. Koo J, Lebwohl A. Psychodermatology: the mind and skin connection. *Am Fam Physician* 2001; 64:1873-1878.
4. Moses S. Pruritic condition. In *Family Practice Notebook* [database online]. 2000; updated June 6, 2004. Available at: www.fpnotebook.com/DER258.htm. Accessed July 23, 2004.
5. Moses S. Pruritus causes. In *Family Practice Notebook* [database online]. 2000; updated June 6, 2004. Available at: www.fpnotebook.com/DER259.htm. Accessed July 23, 2004.

Note: A handout developed by the American Academy of Family Physicians for patients with neurotic excoriations is available to print or photocopy at the following website: www.aafp.org/afp/20011215/neurph.html.

DRUG BRAND NAMES

Cephalexin • Biocef, Keflex, Keftab
Dicloxacillin • Dycill, Dynapen, Pathocil
Doxepin • Adapin, Sinequan