

CLINICAL INQUIRIES

TABLE

Warning signs and symptoms of dyspepsia and GERD that suggest complicated disease or more serious underlying process¹

Dysphagia
 Unexplained weight loss
 History of gastrointestinal bleeding
 Early satiety
 Iron deficiency anemia
 Vomiting
 Odynophagia (sharp substernal pain on swallowing)
 Initial onset of heartburn-like symptoms after the age of 50 years
 History of immunocompromised state
 Anorexia

555 patients with endoscopically proven reflux esophagitis, 154 of whom were over the age of 65. After 8 weeks, rates of esophageal healing among the elderly were 70% for those receiving omeprazole and 29% for those receiving H2RAs (41% difference; 95% confidence interval [CI], 26–55), while the rate of asymptomatic elderly patients was 79% for the omeprazole group and 51% for the H2RA group (28% difference; 95% CI, 12–44).³ Patients treated with omeprazole healed faster than those taking H2RAs, as shown by endoscopy, and more of them experienced symptom relief.

A multicenter, randomized, double-blind trial of GERD maintenance therapy started with an initial open phase in which elderly patients with GERD and documented esophagitis were treated and then had documented resolution of esophagitis by endoscopy after 6 months. The researchers then randomized 105 of these elderly patients to receive treatment with either low-dose (20 mg/d) pantoprazole or placebo for 6 months. Endoscopy was performed after 12 months for all patients, unless indicated sooner. Intention-to-treat analysis showed a disease-free rate of 79.6% (95% CI, 68.3–90.9) in the treatment group, compared with 30.4% (95% CI, 18.3–42.4) in the placebo group (number needed to treat [NNT]=2). Symptom reports concerning the same patients also suggest a marked drop in symptoms that correlated with healing.⁴

A prospective, nonequivalence, before-after study compared efficacy of, and complications from, laparoscopic surgery for symptomatic GERD between younger and older (≥ 65 years) patients. The investigators examined postoperative morbidity and mortality for 359 patients referred for laparoscopic surgery, either a partial or Nissen (full) fundoplication. They excluded those requiring more extensive surgery or repair of paraesophageal hernia. The 42 elderly patients had a higher mean American Society of Anesthesiologists score compared with the younger patients, reflecting higher preoperative comorbidity, but were similar with regard to weight and gender.

Before surgery, investigators performed 24-hour ambulatory pH monitoring. Preoperative exposure times to a pH below 4 (TpH <4) were similar for the younger and older patients (median 14.2% and 13.9%, respectively). Postoperative complication rates were similar for both groups. No deaths occurred. Minor postoperative complications involved 7% of the elderly patients and 6% of the younger group. The 24-hour pH monitoring scores showed improvement at 6 weeks after surgery for both groups, with the median TpH <4 at 1.1% (95% CL, 0.5) in the elderly vs a median of 1.8% (95% CL, 1.9) in the younger patients. At 1 year postoperatively, the values were also similar between the two groups; the median TpH <4 (95% CL) were 1.4% (1.5) in the elderly group and 1.2% (0.6) in the younger patient group.

The results of this study should be interpreted with caution, however. The study design is prone to bias, the patients had relatively low symptom scores at baseline, and sicker patients may have been excluded during the referral process.⁵

Recommendations from others

The Veterans Health Affairs/Department of Defense clinical practice guidelines recommend differentiating GERD (feelings of substernal burning associated with acid regurgitation) from dyspepsia (chronic or recurrent discomfort centered in the upper

FAST TRACK

Based on safety profiles and success in the general population, PPIs as a class are first-line treatment for GERD in the elderly

CONTINUED ON PAGE 258

CLINICAL INQUIRIES

What is the best way to manage GERD symptoms in the elderly?

Brian S. Bacak, MD, Mihir Patel, MD

Rose Family Medicine Residency, University of Colorado Health Sciences Center

Elizabeth Tweed, MLIS

Denison Memorial Library, University of Colorado at Denver and Health Sciences Center

EVIDENCE-BASED ANSWER

No evidence supports one method over another in managing uncomplicated gastroesophageal reflux disease (GERD) for patients aged >65 years. For those with endoscopically documented esophagitis, proton pump inhibitors (PPIs) relieve symptoms faster than histamine H2 receptor antagonists (H2RAs) (strength of recommendation [SOR]: **B**, extrapolation from randomized controlled trials [RCTs]). Treating elderly patients with pantoprazole (Protonix) after resolution of acute esophagitis results in fewer relapses than with placebo (SOR: **B**, double-blind RCT). Limited evidence suggests that such maintenance therapy for prior esophagitis with either H2RAs or PPIs, at half- and full-dose strength,¹ decreases the frequency of relapse (SOR: **B**, extrapolation from uncontrolled clinical trial).

Laparoscopic antireflux surgery for treating

symptomatic GERD among elderly patients without paraesophageal hernia reduces esophageal acidity, with no apparent increase in postoperative morbidity or mortality compared with younger patients (SOR: **C**, nonequivalent before-after study). Upper endoscopy is recommended for elderly patients with alarm symptoms, new-onset GERD, or longstanding disease (SOR: **C**, expert consensus; see **TABLE**).

Elderly patients are at risk for more severe complications from GERD, and their relative discomfort from the disease process is often less than from comparable pathology for younger patients (SOR: **C**, expert consensus). Based on safety profiles and success in the general patient population, PPIs as a class are considered first-line treatment for GERD and esophagitis for the elderly (SOR: **C**, expert consensus).

CLINICAL COMMENTARY

Teasing out serious disease from routine GERD is a challenge in the elderly

Who really needs the invasive workup and expensive medicine, and who needs the simple approach? Elderly patients can attribute symptoms to "just indigestion" when there are other more serious diseases present (ie, angina, severe esophagitis, stricture, cancer). The physician may need a more detailed and patient approach to get the "real" story. Lifestyle changes and over-the-

counter medications will resolve the majority of GERD symptoms. If the simple things don't work or there are warning signs/symptoms, then a further workup is needed. The older patient may then be willing to pay the cost of the long term PPI when they know that there is significant pathology and the potential for long term symptom relief.

Peter Danis, MD
St. John's Mercy Medical Center, St. Louis, Mo

■ Evidence summary

Aggregated data from 2 randomized reflux esophagitis trials conducted in the United Kingdom were analyzed with respect to patient age. Comparison of symptom relief

and esophageal lesion healing showed that elderly patients treated with omeprazole (Prilosec) fared better than those treated with either cimetidine (Tagamet) or ranitidine (Zantac).² The pooled data involved

CONTINUED