



Do probiotics reduce *C diff* risk in hospitalized patients?

A systematic review and meta-analysis says, “Yes,” but that doesn’t necessarily mean they will start appearing on hospital formularies.

PRACTICE CHANGER

Start probiotics within 1 to 2 days of starting antibiotics in hospitalized patients to reduce the risk of *Clostridium difficile* infection.¹

STRENGTH OF RECOMMENDATION

A: Based on a meta-analysis of randomized controlled trials.

Shen NT, Maw A, Tmanova LL, et al. Timely use of probiotics in hospitalized adults prevents *Clostridium difficile* infection: a systematic review with meta-regression analysis. *Gastroenterology*. 2017;152:1889-1900. e9.

ILLUSTRATIVE CASE

A 68-year-old woman is admitted to the hospital with a diagnosis of community-acquired pneumonia. Should you add probiotics to her antibiotic regimen to prevent infection with *Clostridium difficile*?

Clostridium *difficile* infection (CDI) leads to significant morbidity, mortality, and treatment failures. In 2011, it culminated in a cost of \$4.8 billion and 29,000 deaths.^{2,3} Risk factors for infection include antibiotic use, hospitalization, older age, and medical comorbidities.² Probiotics have been proposed as one way to prevent CDI.

While several systematic reviews have demonstrated efficacy for probiotics in the prevention of CDI,⁴⁻⁶ guidelines from the American College of Gastroenterology and the Society for Healthcare Epidemiology of America did not incorporate a recommendation for the use of probiotics in their CDI prevention strategy.^{7,8}

The PLACIDE trial studied the use of probiotics in inpatients ages ≥ 65 years receiving either oral or parenteral antibiotics and found no difference in the incidence of CDI in those who received probiotics vs those who did not.⁹ Even though the PLACIDE trial was the largest, high-quality, randomized controlled trial (RCT) on the use of probiotics to prevent CDI, it had a lower incidence of CDI than was assumed in the power calculations. Additionally, previous systematic reviews did not always follow the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines, and did not focus specifically on hospitalized patients, who are at higher risk for CDI.

Given the conflicting and poor evidence and recommendations, an additional systematic review and meta-analysis was performed following PRISMA guidelines and focusing on studies conducted only on hospitalized adults.

STUDY SUMMARY

Probiotics prevent CDI in hospitalized patients receiving antibiotics

This meta-analysis of 19 RCTs evaluated the efficacy of probiotics for the prevention of CDI in 6261 adult hospitalized patients taking antibiotics. All patients were ≥ 18 years (mean age 68-69 years) and received antibiotics orally, intravenously, or via both routes for any medical indication.

Trials were included if the intervention was for CDI prevention and if the probiotics

Matthew Simpson, MD, MPH; Corey Lyon, DO
University of Colorado
Family Medicine Residency,
Denver

DEPUTY EDITOR
Jennie B. Jarrett,
PharmD, BCPS, MMedEd
University of Illinois
at Chicago

➤ Administration of probiotics to hospitalized patients—particularly when started within 1 to 2 days of initiating antibiotic therapy—can prevent *C diff* infections.

used were 1 or a combination of 4 strains (*Lactobacillus*, *Saccharomyces*, *Bifidobacterium*, *Streptococcus*). Probiotic doses ranged from 4 billion to 900 billion colony-forming u/day and were started from 1 to 7 days after first antibiotic dose. Duration of probiotic use was either fixed at between 14 and 21 days or varied based on the duration of antibiotics (extending 3-14 days after the last antibiotic dose).

Control groups received matching placebo in all trials but 2; those 2 used usual care of no probiotics as the control. Common patient exclusions were pregnancy, immune system compromise, intensive care, a prosthetic heart valve, and pre-existing gastrointestinal disorders.

The risk for CDI was lower in the probiotic group (range 0%-11%) than in the control group (0%-40%) with no heterogeneity ($I^2 = 0.0\%$; $P = .56$) when the data were pooled from all 19 studies (relative risk [RR] = 0.42; 95% confidence interval [CI], 0.30-0.57). The median incidence of CDI in the control groups from all studies was 4%, which yielded a number needed to treat (NNT) of 43 (95% CI, 36-58).

The researchers examined the NNT at varying incidence rates. If the incidence of CDI was 1.2%, the NNT to prevent 1 case of CDI was 144, and if the incidence was 7.4%, the NNT was 23. Compared with control groups, there was a significant reduction in CDI if probiotics were started within 1 to 2 days of antibiotic initiation (RR = 0.32; 95% CI, 0.22-0.48), but not if they were started at 3 to 7 days (RR = 0.70; 95% CI, 0.40-1.2). There was no significant difference in adverse events (ie, cramping, nausea, fever, soft stools, flatulence, taste disturbance) between probiotic and control groups (14% vs 16%; $P = .35$).

WHAT'S NEW

Probiotics provide added benefit if taken sooner rather than later

This high-quality meta-analysis shows that administration of probiotics to hospitalized patients—particularly when started within 1 to 2 days of initiating antibiotic therapy—can prevent CDI.

CAVEATS

Findings do not apply to all patients; specific recommendations are lacking

Findings from this meta-analysis do not apply to patients who have an immunocompromising condition, are pregnant, have a prosthetic heart valve, have a pre-existing gastrointestinal disorder (eg, irritable bowel disease, pancreatitis), or require intensive care. In addition, specific recommendations as to the optimal probiotic species, dose, formulation, and duration of use cannot be made based on this meta-analysis. Lastly, findings from this study do not apply to patients treated with antibiotics in the ambulatory care setting.

CHALLENGES TO IMPLEMENTATION

Lack of “medication” status leads to limited availability in hospitals

The largest barrier to giving probiotics to hospitalized adult patients is the availability of probiotics on local hospital formularies. Probiotics are not technically a medication; they are not regulated or approved by the US Food and Drug Administration and thus, insurance coverage and availability for inpatient use are limited. Lastly, US cost-effectiveness data are lacking, although such data would likely be favorable given the high costs associated with treatment of CDI.

JFP

ACKNOWLEDGMENT

The PURLs Surveillance System was supported in part by Grant Number UL1RR024999 from the National Center For Research Resources, a Clinical Translational Science Award to the University of Chicago. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Center For Research Resources or the National Institutes of Health.

Copyright © 2019. The Family Physicians Inquiries Network. All rights reserved.

References

- Shen NT, Maw A, Tmanova LL, et al. Timely use of probiotics in hospitalized adults prevents *Clostridium difficile* infection: a systematic review with meta-regression analysis. *Gastroenterology*. 2017;152:1889-1900.e9.
- Evans CT, Safdar N. Current trends in the epidemiology and outcomes of *Clostridium difficile* infection. *Clin Infect Dis*. 2015;60(Suppl 2):S66-S71.
- Lessa FC, Winston LG, McDonald LC, et al. Burden of *Clostridium difficile* infection in the United States. *N Engl J Med*. 2015;372:2369-2370.
- Goldenberg JZ, Yap C, Lytvyn L. Probiotics for the prevention of *Clostridium difficile*-associated diarrhea in adults and children. *Cochrane Database Syst Rev*. 2017;12:CD006095.

CONTINUED

5. Lau CS, Chamberlain RS. Probiotics are effective at preventing *Clostridium difficile*-associated diarrhea: a systematic review and meta-analysis. *Int J Gen Med.* 2016;22:27-37.

6. Johnston BC, Goldenberg JZ, Guyatt GH. Probiotics for the prevention of *Clostridium difficile*-associated diarrhea. In response. *Ann Intern Med.* 2013;158:706-707.

7. Surawicz CM, Brandt LJ, Binion DG, et al. Guidelines for diagnosis, treatment, and prevention of *Clostridium difficile* infections. *Am J Gastroenterol.* 2013;108:478-498.

8. Cohen SH, Gerding DN, Johnson S, et al. Clinical practice guidelines for *Clostridium difficile* infection in adults: 2010 update by the Society for Healthcare Epidemiology of America (SHEA) and the Infectious Diseases Society of America (IDSA). *Infect Control Hosp Epidemiol.* 2010;31:431-455.

9. Allen SJ, Wareham K, Wang D, et al. Lactobacilli and bifidobacteria in the prevention of antibiotic-associated diarrhoea and *Clostridium difficile* diarrhoea in older inpatients (PLACIDE): a randomised, double-blind, placebo-controlled, multicentre trial. *Lancet.* 2013;382:1249-1257.

Join the official job board of
 THE JOURNAL OF
**FAMILY
 PRACTICE**

 **MEDJOBNETWORK.com**
 Physician • NP/PA Career Center

**Search 1000s of Jobs and Apply in 1 Click
 And get FREE benefits including...**

- Access to 30+ medical web sites
- Medical news and clinical advice
- E-Alerts and Newsletters on your smart phone
- Online CME and MD-IQ Quizzes
- Coverage of over 200 meetings
- And much more!

Julian Knight
 Digital Sales Director, Classifieds
 Phone: 973-206-2317
 jknight@mdedge.com
 www.medjobnetwork.com