

# Light Therapy Underwhelming for Acne

BY DOUG BRUNK  
San Diego Bureau

CARLSBAD, CALIF. — Some acne patients may respond to various light-based treatments, but most of the time the improvement is modest and short-lived, Dr. E. Victor Ross Jr. said at a symposium on laser and cosmetic surgery sponsored by SkinCare Physicians.

Most light-based therapies for acne “don’t work as well as they should,” said Dr. Ross, director of the Scripps Clinic Skin and Cosmetic Center in La Jolla, Calif. “The bar should be pretty high. If we’re having patients sit under these lights and have these therapies, I think that means long remission periods with one treatment, not five treatments every 2 or 3 weeks, and evidence that there’s some robust compromise of the sebaceous gland.”

Most patients who seek light-based treatments for their acne have been on isotretinoin in the past yet are unwilling to be on systemic retinoids again. “They don’t have severe acne, but they have cyclical moderate acne, and they’re tired and frustrated with the typical” therapies, Dr. Ross said.

Light therapies don’t work well for acne because a photochemical effect should prevail over any photothermal effect to excite endogenous porphyrins produced by *Propionibacterium acnes*. In addition, the only studies to show microscopic damage

to the sebaceous gland have been those with long incubation times, continuous-wave light sources, and red light only. This regimen caused epidermal damage in every case, he noted.

“It’s unlikely that one will achieve long-term and profound sebaceous gland compromise with short aminolevulinic acid times, either with pulsed light sources or with continuous-wave light sources,” he said. “Right now we have to be honest with ourselves and say the prescription pad is still pretty darned good. ... I’ve tried

these [light] therapies, but I’m not ready to say these are a home run or even a triple or double right now.”

He added that many patients who undergo light treatment for their acne fail what he calls the “come back” test. “They don’t come back. If they don’t come back, [we assume] they’re probably better, but I suspect that most of them are not better, they’re worse,” he said.

Dr. Ross disclosed that he has research relationships with Palomar, Cutera, and Laserscope. ■

## VERBATIM

*‘I always make it very clear ... that I am a dermatologist and not a plastic surgeon. I take pride in the specialty and promote dermatologists’ unique perspective and skills.’*

Dr. Greg S. Morganroth, on the surgical consent process, p. 76

## Anticoagulants Not Precluded Before Surgery

SAN DIEGO — Dermatologists should ask patients detailed questions about what medications they are taking before performing a procedure, Dr. Rainer E. Sachse said at a meeting sponsored by the American Society for Mohs Surgery.

Almost half of patients who undergo a dermatologic procedure are taking some type of anticoagulant, according to a recent survey. However, this scenario may not preclude surgery, Dr. Sachse said.

Patients may be taking an alternative medicine that has anticoagulant properties, such as garlic, fish oil, niacin, or vitamin E. Similarly, many patients routinely take aspirin each day, noted Dr. Sachse, a facial plastic surgeon who practices Mohs surgery in Fort Lauderdale, Fla.

If a patient can stop the anticoagulant, Dr. Sachse asks them to do so for at least 1 week before the procedure. But because of the possible risk involved in stopping that therapy, he has the patient consult with the doctor who prescribed the medication.

For patients who are taking warfarin (Coumadin), dermatologists should know the international normalized ratio before performing the procedure. A patient on any type of anticoagulant should keep the pressure dressing on longer and avoid a lot of activity afterward to prevent bruising, other physicians at the meeting said.

—Timothy F. Kirn



## Safety that’s reassuring for everyone

For children and adults,  
Cloderm® is the mid-potency topical steroid  
you can prescribe with confidence.

- Clinically proven to preferentially remain in the epidermis to enhance safety<sup>1</sup>
- Rapid response—as early as Day 4<sup>1</sup>

**Class C**  
corticosteroid:  
no significant cross-reactivity<sup>2,3</sup>

- The most common adverse events with Cloderm include dryness, irritation, folliculitis, acneiform eruptions, and burning. Cloderm is contraindicated in patients who are hypersensitive to any of the ingredients of this product. As with all topical corticosteroids, systemic absorption can produce reversible HPA-axis suppression.

Please see full prescribing information on reverse side of page.

References: 1. Data on file, Healthpoint, Ltd. 2. Jacob SE, Steele T. Corticosteroid classes: A quick reference guide including patch test substances and cross-reactivity. *J Am Acad Dermatol*. 2006;54(4):723-727. 3. Matura M, Goossens A. Contact allergy to corticosteroids. *Allergy*. 2000;55:698-704.

Cloderm is a registered trademark of Healthpoint, Ltd.  
©2006 CORIA Laboratories, Ltd. A DFB Company. CL-20016

**CORIA**  
LABORATORIES, LTD.  
A DFB COMPANY  
www.corialabs.com

**Cloderm**®  
(clocortolone pivalate)  
Cream, 0.1%

