

Interview Software Evaluates Autism Symptoms

BY HEIDI SPLETE
Senior Writer

The assessment of children with autism spectrum disorders is going high tech.

The Developmental, Dimensional and Diagnostic Interview, known as 3di, is a computerized program that analyzes autistic symptoms in clinical and normal populations, said David Skuse, M.D., of University College London and his colleagues.

Current standardized interviews, including the National Institute of Mental Health Diagnostic Interview Schedule for Children, fail to measure autistic symptoms, the researchers noted.

Although nothing substitutes for a careful interview by someone who knows the child well, the 3di is a hybrid of structured and semistructured interview methods (J. Am. Acad. Child Adolesc. Psychiatry 2004;43:548-58).

The program is meant primarily to

evaluate autistic traits in children with normal-range abilities, but it includes features to assess children with moderate or severe mental retardation as well, the researchers said.

In terms of concurrent validity, the 3di was nearly perfect in determining whether children had any autism diagnosis: Among 60 cases and 60 comparison children (mean age 11 years), the 3di diagnosed only one comparison patient with generalized autism disorder.

In a second approach to concurrent validity involving 60 symptomatic children who were referred to psychiatric services, 27 had clinician-diagnosed autism spectrum disorders. Based on how well the children met the International Statistical Classification of Diseases, the 3di program diagnosed 29 of the children with a significant autistic disorder. In addition, the 3di's positive predictive power was 0.93 and negative predictive power was 0.91 in a discriminant validity study of the program's ability to distinguish autism spectrum disorders from nonautistic conditions.

Dr. Skuse is a stockholder in IXDX Ltd., which owns the rights to the interview software and to the dissemination of the technology.

Study Supports Autism's Link To Immunity

BUDAPEST, HUNGARY — Children with autism spectrum disorders appear to have immune responses different from those of healthy children, according to data presented at the 4th International Congress on Autoimmunity.

"There is evidence now for an immune dysregulation in children with autism, compared with children in the general population," said Paul Ashwood, Ph.D., of the University of California, Davis.

In a study of 31 children ASD and 19 typically developing control children aged 2-5 years, the children with ASD had abnormal levels of several cytokines in response to stimulation with three antigens, compared with the control children.

The study adds weight to the idea that autism has an immune component. There are several previous reports of both increased autoimmunity and immune response deficits in children with autism spectrum disorders (ASD). "However, a lot of these reports are conflicting, and there is no consensus so far," Dr. Ashwood said.

Dr. Ashwood and colleagues isolated and stimulated peripheral blood mononuclear cells for 48 hours with phytohemagglutinin, lipopolysaccharide, and vaccine antigens. Analysis was performed for 18 cytokines. At baseline, cytokine levels were similar in the children with ASD and the control children.

Following stimulation with phytohemagglutinin, the children with ASD had statistically significantly lower levels of IL-2, IL-6, IL-10, and IL-12 than the control children. There was also a trend toward higher levels of IL-13 and granulocyte macrophage-colony stimulating factor, compared with the control children.

—Kerri Wachter

EQUETRO™ (carbamazepine) extended-release capsules
100 mg, 200 mg and 300 mg

Rx only

Brief Summary Prescribing Information

WARNING: APLASTIC ANEMIA AND AGRANULOCYTOSIS HAVE BEEN REPORTED IN ASSOCIATION WITH THE USE OF CARBAMAZEPINE. DATA FROM A POPULATION-BASED CASE-CONTROL STUDY DEMONSTRATE THAT THE RISK OF DEVELOPING THESE REACTIONS IS 5-8 TIMES GREATER THAN IN THE GENERAL POPULATION. HOWEVER, THE OVERALL RISK OF THESE REACTIONS IN THE UNTREATED GENERAL POPULATION IS LOW. APPROXIMATELY SIX PATIENTS PER ONE MILLION POPULATION PER YEAR FOR AGRANULOCYTOSIS AND TWO PATIENTS PER ONE MILLION POPULATION PER YEAR FOR APLASTIC ANEMIA. ALTHOUGH REPORTS OF TRANSIENT OR PERSISTENT DECREASED PLATELET OR WHITE BLOOD CELL COUNTS ARE NOT UNCOMMON IN ASSOCIATION WITH THE USE OF CARBAMAZEPINE, DATA ARE NOT AVAILABLE TO ESTIMATE ACCURATELY THEIR INCIDENCE OR OUTCOME. HOWEVER, THE VAST MAJORITY OF THE CASES OF LEUKOPENIA HAVE NOT PROGRESSED TO THE MORE SERIOUS CONDITIONS OF APLASTIC ANEMIA OR AGRANULOCYTOSIS. BECAUSE OF THE VERY LOW INCIDENCE OF AGRANULOCYTOSIS AND APLASTIC ANEMIA, THE VAST MAJORITY OF MINOR HEMATOLOGIC CHANGES OBSERVED IN MONITORING OF PATIENTS ON CARBAMAZEPINE ARE UNLIKELY TO SIGNAL THE OCCURRENCE OF EITHER ABNORMALITY. NONETHELESS, COMPLETE PRETREATMENT HEMATOLOGICAL TESTING SHOULD BE OBTAINED AS A BASELINE. IF A PATIENT IN THE COURSE OF TREATMENT EXHIBITS LOW OR DECREASED WHITE BLOOD CELL OR PLATELET COUNTS, THE PATIENT SHOULD BE MONITORED CLOSELY. DISCONTINUATION OF THE DRUG SHOULD BE CONSIDERED IF ANY EVIDENCE OF SIGNIFICANT BONE MARROW DEPRESSION DEVELOPS.

Before prescribing EQUETRO™, the physician should be thoroughly familiar with the details of the full prescribing information, particularly regarding use with other drugs, especially those which accentuate toxicity potential.

INDICATIONS AND USAGE

EQUETRO™ is indicated for the treatment of acute manic and mixed episodes associated with Bipolar I Disorder. The efficacy of EQUETRO™ in acute mania was established in 2 placebo-controlled, double-blind, 3-week studies in patients meeting DSM-IV criteria for Bipolar I Disorder who currently displayed an acute manic or mixed episode. The effectiveness of EQUETRO™ for longer-term use and for prophylactic use in mania has not been systematically evaluated in controlled clinical trials. Therefore, physicians who elect to use EQUETRO™ for extended periods should periodically re-evaluate the long-term risks and benefits of the drug for the individual patient (see **DOSE AND ADMINISTRATION**).

CONTRAINDICATIONS

Carbamazepine should not be used in patients with a history of previous bone marrow depression, hypersensitivity to the drug, or known sensitivity to any of the tricyclic compounds, such as amitriptyline, desipramine, imipramine, protriptyline and nortriptyline. Likewise, on theoretical grounds its use with monoamine oxidase inhibitors is not recommended. Before administration of carbamazepine, MAO inhibitors should be discontinued for a minimum of 14 days, or longer if the clinical situation permits.

WARNINGS

Patients should be made aware that EQUETRO™ contains carbamazepine and should not be used in combination with any other medications containing carbamazepine.

Use in Pregnancy

Carbamazepine can cause fetal harm when administered to a pregnant woman. Epidemiological data suggest that there may be an association between the use of carbamazepine during pregnancy and congenital malformations, including spina bifida. The prescribing physician will wish to weigh the benefits of therapy against the risks in treating or counseling women of childbearing potential. If this drug is used during pregnancy, or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to the fetus. Retrospective case reviews suggest that, compared with monotherapy, there may be a higher prevalence of teratogenic effects associated with the use of anticonvulsants in combination therapy. In humans, transplacental passage of carbamazepine is rapid (30-60 minutes), and the drug is accumulated in the fetal tissues with higher levels found in liver and kidney than in brain and lung. Carbamazepine has been shown to have adverse effects in reproduction studies in rats when given orally in dosages 10-25 times a human daily dosage of 1200 mg on a mg/kg basis or 1.5-4 times the human daily dosage on a mg/m² basis. In rat teratology studies, 2 of 135 offspring showed kinked ribs at 250 mg/kg and 4 of 119 offspring at 650 mg/kg showed other anomalies (cleft palate, 1; talipes, 1; anophthalmos, 2). In reproduction studies in rats, nursing offspring demonstrated a lack of weight gain and an unkempt appearance at a maternal dosage level of 200 mg/kg.

Tests to detect defects using current accepted procedures should be considered a part of routine prenatal care in childbearing women receiving carbamazepine.

General

Patients with a history of adverse hematologic reaction to any drug may be particularly at risk. Severe dermatologic reactions, including toxic epidermal necrolysis (Lyell's syndrome) and Stevens-Johnson syndrome have been reported with carbamazepine. These reactions have been extremely rare. However, a few fatalities have been reported.

In patients with seizure disorder, carbamazepine should not be discontinued abruptly because of the strong possibility of precipitating status epilepticus with attendant hypoxia and threat to life. Carbamazepine has shown mild anticholinergic activity; therefore, patients with increased intraocular pressure should be closely observed during therapy. Because of the relationship of the drug to other tricyclic compounds, the possibility of activation of a latent psychosis and, in elderly patients, of confusion or agitation should be considered. Co-administration of carbamazepine and delavirdine may lead to loss of virologic response and possible resistance to the class of non-nucleoside reverse transcriptase inhibitors.

PRECAUTIONS

General

Before initiating therapy, a detailed history and physical examination should be made. Therapy should be prescribed after critical benefit-risk appraisal in patients with a history of cardiac, hepatic, or renal damage; adverse hematologic reaction to other drugs; or interrupted courses of therapy with carbamazepine.

Suicide: The possibility of suicide attempt is inherent in Bipolar Disorder and close supervision of high risk patients should accompany drug therapy. Prescriptions for EQUETRO™ should be written for the smallest quantity consistent with good patient management in order to reduce the risk of overdose.

Information for Patients

Patients should be made aware of the early toxic signs and symptoms of a potential hematologic problem, such as fever, sore throat, rash, ulcers in the mouth, easy bruising, petechial or purpuric hemorrhage, and should be advised to report to the physician immediately if any such signs or symptoms appear. Since dizziness and drowsiness may occur, patients should be cautioned about the hazards of operating machinery or automobiles or engaging in other potentially dangerous tasks. If necessary, the EQUETRO™ capsules can be opened and the contents sprinkled over food, such as a teaspoon of applesauce or other similar food products. EQUETRO™ capsules or their contents should not be crushed or chewed. EQUETRO™ may interact with some drugs. Therefore, patients should be advised to report to their doctors the use of any other prescription or non-prescription medication or herbal products.

Laboratory Tests

Complete pretreatment blood counts, including platelets and possibly reticulocytes and serum iron, should be obtained as a baseline. If a patient in the course of treatment exhibits low or decreased white blood cell or platelet counts, the patient should be monitored closely. Discontinuation of the drug should be considered if any evidence of significant bone marrow depression develops.

Baseline and periodic evaluations of liver function, particularly in patients with a history of liver disease, must be performed during treatment with this drug since liver damage may occur. The drug should be discontinued immediately in cases of aggravated liver dysfunction or active liver disease.

Baseline and periodic eye examinations, including slit-lamp, funduscopy, and tonometry, are recommended since many phenothiazines and related drugs have been shown to cause eye changes.

Baseline and periodic complete urinalysis and BUN determinations are recommended for patients treated with this agent because of observed renal dysfunction.

Increases in total cholesterol, LDL and HDL have been observed in some patients taking anticonvulsants. Therefore, periodic evaluation of these parameters is also recommended.

Monitoring of blood levels (please see full prescribing information) may be useful for verification of drug compliance, assessing safety and determining the cause of toxicity including when more than one medication is being used.

Thyroid function tests have been reported to show decreased values with carbamazepine administered alone. Hyponatremia has been reported in association with carbamazepine use, either alone or in combination with other drugs. Interference with some pregnancy tests has been reported.

Drug Interactions

Clinically meaningful drug interactions have occurred with concomitant medications and include, but are not limited to the following:

Agents Highly Bound to Plasma Protein: Carbamazepine is not highly bound to plasma proteins; therefore, administration of EQUETRO™ to a patient taking another drug that is highly protein bound should not cause increased free concentrations of the other drug.

Agents that Inhibit Cytochrome P450 Isoenzymes and/or Epoxide Hydrolase: Carbamazepine is metabolized mainly by cytochrome P450 (CYP) 3A4 to the active carbamazepine 10,11-epoxide, which is further metabolized to the trans-diol by epoxide hydrolase. Therefore, the potential exists for interaction between carbamazepine and any agent that inhibits CYP3A4 and/or epoxide hydrolase. CYP3A4 inhibitors have been found, or are expected, to increase plasma levels of EQUETRO™. Commonly used agents that inhibit CYP3A4 are: azole antifungals (such as ketoconazole and itraconazole, calcium channel blockers (such as diltiazem and verapamil), macrolide antibiotics (such as erythromycin, clarithromycin, and troleandomycin), grapefruit juice, and other drugs. Please see full prescribing information.

Thus, if a patient has been titrated to a stable dosage of EQUETRO™, and then begins a course of treatment with one of these CYP3A4 or epoxide hydrolase inhibitors, it is reasonable to expect that a dose reduction for EQUETRO™ may be necessary.

Agents that Induce Cytochrome P450 Isoenzymes: Carbamazepine is metabolized by CYP3A4. Therefore, the potential exists for interaction between carbamazepine and any agent that induces CYP3A4. CYP3A4 inducers have been found, or are expected, to decrease plasma levels of EQUETRO™. Commonly used agents that induce CYP3A4 are: phenytoin, primidone, theophylline, anticancer agents, and other drugs. Please see full prescribing information. Thus, if a patient has been titrated to a stable dosage on EQUETRO™, and then begins a course of treatment with one of these CYP3A4 inducers, it is reasonable to expect that a dose increase for EQUETRO™ may be necessary.

Agents with Decreased Levels in the Presence of Carbamazepine due to Induction of Cytochrome P450 Enzymes: Carbamazepine is known to induce CYP1A2 and CYP3A4. Therefore, the potential exists for interaction between carbamazepine and any agent metabolized by one (or more) of these enzymes. These agents have been found, or are expected to have decreased plasma levels in the presence of EQUETRO™ due to induction of CYP enzymes. Commonly used agents that induce CYP enzymes are: acetaminophen, benzodiazepines (such as alprazolam, diazepam, lorazepam, midazolam, and triazolam), protease inhibitors, oral contraceptives, antidepressants (tricyclics and SSRIs), phenytoin, and other drugs. Please see full prescribing information. Break through bleeding has been reported among patients receiving concomitant oral contraceptives and their reliability may be adversely affected.

Warfarin's anticoagulant effect can be reduced in the presence of carbamazepine.

Thus, if a patient has been titrated to a stable dosage on one of the agents in this category, and then begins a course of treatment with EQUETRO™, it is reasonable to expect that a dose increase for the concomitant agent may be necessary.

Agents with Increased Levels in the Presence of Carbamazepine: EQUETRO™ increases the plasma levels of clomipramine HCl and primidone.

Thus, if a patient has been titrated to a stable dosage on one of the agents in this category, and then begins a course of treatment with EQUETRO™, it is reasonable to expect that a dose decrease for the concomitant agent may be necessary.

Phenytoin has been reported to decrease or increase in the presence of carbamazepine. Careful monitoring of phenytoin plasma levels following co-medication with carbamazepine is advised.

Pharmacological/Pharmacodynamic Interactions with Carbamazepine

Concomitant administration of carbamazepine and lithium may increase the risk of neurotoxic side effects.

Because the anticonvulsant properties of carbamazepine, EQUETRO™ may reduce the thyrotoxic function as has been reported with other anticonvulsants. Additionally, anti-malarial drugs, such as chloroquine and mefloquine, may antagonize the activity of carbamazepine.

Thus if a patient has been titrated to a stable dosage on one of the agents in this category, and then begins a course of treatment with EQUETRO™, it is reasonable to expect that a dose adjustment may be necessary.

Because of its primary CNS effect, caution should be used when EQUETRO™ is taken with other centrally acting drug agents.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Administration of carbamazepine to Sprague-Dawley rats for two years in the diet at doses of 25, 75, and 250 mg/kg/day (low dose approximately 0.2 times the human daily dose of 1200 mg on a mg/m² basis), resulted in a dose-related increase in the incidence of hepatocellular tumors in females and of benign interstitial cell adenomas in the testes of males.

Carbamazepine must, therefore, be considered to be carcinogenic in Sprague-Dawley rats. Bacterial and mammalian mutagenicity studies using carbamazepine produced negative results. The significance of these findings relative to the use of carbamazepine in humans is, at present, unknown.

Use in Pregnancy

Pregnancy Category D (See WARNINGS).

Labor and Delivery

The effect of carbamazepine on human labor and delivery is unknown.

Nursing Mothers

Carbamazepine and its epoxide metabolite are transferred to breast milk and during lactation. Because of the potential for serious adverse reactions in nursing infants from carbamazepine, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use

The safety and effectiveness of EQUETRO™ in pediatric and adolescent patients have not been established.

Geriatric Use

No systematic studies in geriatric patients have been conducted.

ADVERSE REACTIONS

General: The most severe adverse reactions previously observed with carbamazepine were reported in the hemopoietic system (see **BOX WARNING**), the skin, and the cardiovascular system.

The most frequently observed adverse reactions, particularly during the initial phases of therapy, are dizziness, drowsiness, unsteadiness, nausea, and vomiting. To minimize the possibility of such reactions, therapy should be initiated at the lowest dosage recommended.

The most commonly observed adverse experiences (5% and at least twice placebo) seen in association with the use of EQUETRO™ (400 to 1600 mg/day, dose adjusted in 200 mg daily increments in week 1 in Bipolar I Disorder) in the double-blind, placebo-controlled trials of 3 weeks' duration are: dizziness, somnolence, nausea, vomiting, ataxia, pruritus, dry mouth, amblyopia, and speech disorder.

EQUETRO™ and placebo-treated patients from the two double-blind, placebo-controlled studies were enrolled in a 6-month open-label study. The most common adverse events with an incidence of 5% or more are: headache, dizziness, rash, infection, pain, somnolence, diarrhea, dyspepsia, nausea, asthenia, amnesia, accidental injury, anxiety, depression¹, malaise, depressive reaction, chest pain, back pain, constipation, ataxia, and pruritus.

¹Amnesia includes poor memory, forgetful and memory disturbance.

²Depression includes suicidal ideation.

Other significant adverse events seen in less than 5% of patients include: suicide attempt, manic reaction, insomnia, nervousness, depersonalization and extrapyramidal symptoms, infections (fungal, viral, bacterial), pharyngitis, rhinitis, sinusitis, bronchitis, urinary tract infection, leukopenia and lymphadenopathy liver function tests abnormal, edema, peripheral edema, allergic reaction, photosensitivity reaction, alopecia, diplopia and ear pain.

Hemopoietic System: Aplastic anemia, agranulocytosis, pancytopenia, bone marrow depression, thrombocytopenia, leukopenia, leukocytosis, eosinophilia, acute intermittent porphyria.

Skin: Pruritic and erythematous rashes, urticaria, toxic epidermal necrolysis (Lyell's syndrome) (see **WARNINGS**), Stevens-Johnson syndrome (see **WARNINGS**), photosensitivity reactions, alterations in skin pigmentation, exfoliative dermatitis, erythema multiforme and nodosum, purpura, aggravation of disseminated lupus erythematosus, alopecia, and diaphoresis. In certain cases, discontinuation of therapy may be necessary. Isolated cases of hirsutism have been reported, but a causal relationship is not clear.

Cardiovascular System: Congestive heart failure, edema, aggravation of hypertension, hypotension, syncope and collapse, aggravation of coronary artery disease, arrhythmias and AV block, thrombophlebitis, thromboembolism, and adenopathy or lymphadenopathy. Some of these cardiovascular complications have resulted in fatalities. Myocardial infarction has been associated with other tricyclic compounds.

Liver: Abnormalities in liver function tests, cholestatic and hepatocellular jaundice, hepatitis.

Respiratory System: Pulmonary hypersensitivity characterized by fever, dyspnea, pneumonitis, or pneumonia.

Genitourinary System: Urinary frequency, acute urinary retention, oliguria with elevated blood pressure, azotemia, renal failure, and impotence. Albuminuria, glycosuria, elevated BUN, and microscopic deposits in the urine have also been reported. Testicular atrophy occurred in rats receiving carbamazepine orally from 4-52 weeks at dosage levels of 50-400 mg/kg/day. Additionally, rats receiving carbamazepine in the diet for 2 years at dosage levels of 25, 75, and 250 mg/kg/day had a dose-related incidence of testicular atrophy and aspermatogenesis. In dogs, it produced a brownish discoloration, presumably a metabolite, in the urinary bladder at dosage levels of 50 mg/kg/day and higher. Relevance of these findings to humans is unknown.

Nervous System: Dizziness, drowsiness, disturbances of coordination, confusion, headache, fatigue, blurred vision, visual hallucinations, transient diplopia, oculomotor disturbances, nystagmus, speech disturbances, abnormal involuntary movements, peripheral neuritis and paresthesias, depression with agitation, talkativeness, tinnitus, and hyperacusis.

There have been reports of associated paralysis and other symptoms of cerebral arterial insufficiency, but the exact relationship of these reactions to the drug has not been established.

Isolated cases of neuroleptic malignant syndrome have been reported with concomitant use of psychotropic drugs.

Digestive System: Nausea, vomiting, gastric distress and abdominal pain, diarrhea, constipation, anorexia, and dryness of the mouth and pharynx, including glossitis and stomatitis.

Eyes: Scattered punctate cortical lens opacities, as well as conjunctivitis, have been reported. Although a direct causal relationship has not been established, many phenothiazines and related drugs have been shown to cause eye changes.

Musculoskeletal System: Aching joints and muscles, and leg cramps.

Metabolism: Fever and chills, inappropriate antidiuretic hormone (ADH) secretion syndrome has been reported. Cases of frank water intoxication, with decreased serum sodium (hyponatremia) and confusion have been reported in association with carbamazepine use (see **PRECAUTIONS**, Laboratory Tests). Decreased levels of plasma calcium have been reported.

Other: Isolated cases of a lupus erythematosus-like syndrome have been reported. There have been occasional reports of elevated levels of cholesterol, HDL cholesterol, and triglycerides in patients taking anticonvulsants. A case of aseptic meningitis, accompanied by myoclonus and peripheral eosinophilia, has been reported in a patient taking carbamazepine in combination with other medications. The patient was successfully dehydrated, and the meningitis reappeared upon rechallenge with carbamazepine.

419 1207 001 (rev. 12/2004) Manufactured for: **Shire US Inc.**
725 Chesterbrook Blvd., Wayne, PA 19087-5637
1-800-828-2088, Made in U.S.A.
©2005 Shire US Inc.

Shire