

## Hashimoto's Raises Risk of Thyroid Cancer

BY MITCHEL L. ZOLER  
Philadelphia Bureau

PALM BEACH, FLA. — Patients with Hashimoto's thyroiditis had a threefold increased risk of also having a well-differentiated thyroid cancer, compared with other patients undergoing thyroid resection, based on a review of 802 patients treated at one center.

The review also found no link between Hashimoto's thyroiditis (HT) and thyroid lymphoma. Both findings diverge from reports by other groups, which documented a high lymphoma incidence but no increased rate of papillary or follicular thyroid cancers. Dr. B. Mark Evers said at the annual meeting of the Southern Surgical Association.

"There is an association between the inflammation of Hashimoto's thyroiditis and well-differentiated cancers, as in the stomach and colon," said Dr. Evers, professor of surgery at the University of Texas, Galveston. "I'm not sure why we saw more [of an association] than the published literature. This was a carefully done study."

"This is an important observation. It's discordant with most of the published literature. If true, it would have broad ramifications," commented Dr. Robert Udelsman, professor and chairman of surgery at Yale University in New Haven, Conn.

Dr. Evers and associates reviewed all patients who had a thyroid resection at his institution during 1987-2002. The 802 patients included 155 patients with a well-differentiated thyroid cancer only, 52 patients with HT only, 43 with both thyroid cancer and HT, and 552 patients who underwent a resection for another reason, often for laryngectomy. Papillary cancers were 88% of all thyroid cancers, and follicular were another 7%.

Thyroid cancer was about three times more common among patients with Hashimoto's thyroiditis, compared with those in the series without HT, "a strong link between inflammation and thyroid cancer," said Dr. Shawn D. Larson, a surgeon at the University of Texas, Galveston, and a collaborator on the study.

Further analysis showed no association between HT or thyroid cancer and age, gender, race, tumor size, or tumor aggressiveness based on extent of lymph-node involvement and tumor stage.

The demographic and clinical profile of the thyroid cancer patients treated in the Galveston study was compared with more than 22,000 thyroid cancer patients from a database kept by the National Cancer Institute. It showed the Galveston patients were typical, compared with all U.S. patients.

Immunopathology studies of fixed thyroid specimens from randomly selected patients in all four subgroups showed that two proteins involved in cell growth and inflammatory responses were elevated to similar levels in patients with Hashimoto's thyroiditis and those with thyroid cancers, and added support for the hypothesis that the two clinical conditions are linked. ■

## Teen Insulin Resistance Unidentified

BY PATRICE WENDLING  
Chicago Bureau

TUCSON, ARIZ. — There are 1.2 million previously unidentified normal-weight adolescents nationally who may be at increased risk of insulin resistance, Dr. Ann Rodden, said at the annual meeting of the North American Primary Care Research Group.

Adolescents with a body mass index (BMI) in the 75th to 84.9th percentile and

those who have low levels of physical activity were at increased risk for insulin resistance, according to data obtained in a secondary analysis of the National Health and Nutrition Examination Survey (NHANES) during 1999-2002.

Prevalence estimates suggest that more than 8.5 million American adolescents have insulin resistance. Of these, more than 1.2 million are in the 75th to 84.9th BMI percentile. The American Diabetes Association considers adolescents with a

BMI at or above the 85th percentile to be at risk for insulin resistance, said Dr. Rodden, department of family medicine, Medical University of South Carolina, Charleston.

"There is a population of adolescents that right now we do not consider to be at risk of insulin resistance and that we should be looking at in addition to those already identified," she said.

The analysis was based on a nationally representative sample of 1,806 non-

**FIRST**  
IN A NOVEL  
CLASS OF  
**SLEEP**  
AGENTS

Rozerem is indicated for the treatment of insomnia characterized by difficulty with sleep onset. Rozerem can be prescribed for long-term use. Rozerem should not be used in patients with hypersensitivity to any components of the formulation, severe hepatic impairment, or in combination with fluvoxamine. Failure of insomnia to remit after a reasonable period of time should be medically evaluated, as this may be the result of an unrecognized underlying medical disorder. Hypnotics should be administered with caution to patients exhibiting signs and symptoms of depression. Rozerem has not been studied in patients with severe sleep apnea, severe COPD, or in children or adolescents. The effects in these populations are unknown. Avoid taking Rozerem with alcohol. Rozerem has been associated with decreased testosterone levels and increased prolactin levels. Health professionals should be mindful of any unexplained symptoms possibly associated with such changes in these hormone levels. Rozerem should not be taken with or immediately after a high-fat meal. Rozerem should be taken within 30 minutes before going to bed and activities confined to preparing for bed. The most common adverse events seen with Rozerem that had at least a 2% incidence difference from placebo were somnolence, dizziness, and fatigue.

Please see adjacent Brief Summary of Prescribing Information.