

Telbivudine Tops Lamivudine for HBV at 2 Years

BY DIANA MAHONEY
New England Bureau

BOSTON — Telbivudine showed greater efficacy than lamivudine did during 2 years of treatment for chronic hepatitis B infection, according to the most recent results from the international GLOBE trial.

In the phase III trial, telbivudine (Tyze-ka), which recently received Food and Drug Administration approval for the treatment of hepatitis B virus (HBV) infection, sur-

passed the older antiviral on all key measures of virologic and clinical efficacy.

These results were found in both HBV e-antigen (HBeAg)-positive and HBeAg-negative patients enrolled in the study of 1,367 patients, Dr. Ching-Lung Lai said at the annual meeting of the American Association for the Study of Liver Diseases.

The international, multicenter investigation was partially funded by Idenix Pharmaceuticals, the manufacturer—with Novartis Pharma AG—of telbivudine. The

trial was designed to compare the efficacy of telbivudine at 600 mg/day to that of lamivudine (Epivir) at 100 mg/day in patients who were HB s-antigen positive at enrollment and had baseline HBV DNA greater than 6 log₁₀ copies/mL, alanine aminotransferase (ALT) levels between 1.3 and 10 times the upper limit of normal, and compensated liver disease.

The study's primary efficacy end point was therapeutic response, based on a combination of viral suppression (serum HBV

DNA below 100,000 copies/mL) and either improved liver disease markers (ALT normalization) or loss of detectable HBeAg.

The first-year results from the trial, presented last year, "demonstrated that telbivudine was superior to lamivudine on all measures of direct antiviral efficacy and on several key clinical efficacy measures," said Dr. Lai, professor of medicine and hepatology at the University of Hong Kong. "The new data indicate the efficacy levels were sustained over 104 weeks."

After 2 years of treatment, analysis showed that HBeAg-positive and -negative patients randomized to telbivudine therapy had significantly greater HBV DNA reductions (−5.7 log₁₀ and −5.0 log₁₀, respectively) than did HBeAg-positive and -negative patients on lamivudine (−4.4 log₁₀ and −4.2 log₁₀, respectively).

Similarly, significantly more patients receiving telbivudine achieved clearance of detectable HBV DNA. In HBeAg-positive patients, "telbivudine treatment led to loss of detectable HBV DNA in 56% of patients, compared to 39% with lamivudine," said Dr. Lai. Among HBeAg-negative patients, 82% of telbivudine patients achieved undetectable HBV DNA levels, compared with 57% of lamivudine patients, he said.

Patients who had achieved HBV DNA clearance by study week 24 had the best overall response rates in terms of seroconversion, ALT normalization, and reduced therapeutic resistance, Dr. Lai noted.

Treatment failure was significantly less common with telbivudine than with lamivudine in both HBeAg-positive and -negative patients. Among HBeAg-positive patients, antiviral treatment failed in 4% of telbivudine patients, compared with 12% of lamivudine patients. In HBeAg-negative patients, there were no treatment failures in telbivudine patients, compared with 3% of patients taking lamivudine, he said.

Drug resistance occurred in 17.8% of HBeAg-positive patients on telbivudine and 30.1% of HBeAg-positive lamivudine patients during the 2-year study period; resistance was defined in the GLOBE study as HBV DNA return to greater than 5 log₁₀ or to within 1 log₁₀ of baseline. Among the HBeAg-negative patients, 7.3% of those on telbivudine and 16.6% of those on lamivudine developed resistance, Dr. Lai said.

Viral resistance, defined as rebound of HBV DNA after initial suppression to 1 log₁₀ increase above nadir, was found in 21.6% of telbivudine patients and 35.0% of lamivudine patients in the HBeAg-positive group, as well as 8.6% of telbivudine patients and 21.9% of lamivudine patients in the HBeAg-negative group, he said.

Telbivudine was associated with fewer and less severe resistance-associated elevations of serum ALT levels, which occurred in 2.8% of patients, compared with 8.4% of the lamivudine group, Dr. Lai reported.

Both study drugs were generally well tolerated and had similar patterns of clinical adverse events, Dr. Lai said. Grade 3/4 creatine kinase elevations were more commonly associated with telbivudine, occurring in 13% of patients, compared with 4% of lamivudine patients, he noted.

Dr. Lai said he is a scientific adviser to Idenix Pharmaceuticals. ■

Tetanus Toxoid, Reduced Diphtheria Toxoid and Acellular Pertussis Vaccine Adsorbed ADACEL™

Brief Summary: Please see package insert for full prescribing information

INDICATIONS AND USAGE ADACEL vaccine is indicated for active booster immunization for the prevention of tetanus, diphtheria and pertussis as a single dose in persons 11 through 64 years of age. The use of ADACEL vaccine as a primary series, or to complete the primary series, has not been studied. As with any vaccine, ADACEL vaccine may not protect 100% of vaccinated individuals.

CONTRAINDICATIONS Known systemic hypersensitivity to any component of ADACEL vaccine or a life-threatening reaction after previous administration of the vaccine or a vaccine containing the same substances are contraindications to vaccination with ADACEL vaccine. Because of uncertainty as to which component of the vaccine may be responsible, additional vaccinations with the diphtheria, tetanus or pertussis components should not be administered. Alternatively, such individuals may be referred to an allergist for evaluation if further immunizations are to be considered. The following events are contraindications to administration of any pertussis containing vaccine: (1)

- Encephalopathy within 7 days of a previous dose of pertussis containing vaccine not attributable to another identifiable cause.
- Progressive neurological disorder, uncontrolled epilepsy, or progressive encephalopathy. Pertussis vaccine should not be administered to individuals with these conditions until a treatment regimen has been established, the condition has stabilized, and the benefit clearly outweighs the risk.

ADACEL vaccine is not contraindicated for use in individuals with HIV infection. (1)

WARNINGS Because intramuscular injection can cause injection site hematoma, ADACEL vaccine should not be given to persons with any bleeding disorder, such as hemophilia or thrombocytopenia, or to persons on anticoagulant therapy unless the potential benefits clearly outweigh the risk of administration. If the decision is made to administer ADACEL vaccine in such persons, it should be given with caution, with steps taken to avoid the risk of hematoma formation following injection. (1) If any of the following events occurred in temporal relation to previous receipt of a vaccine containing a whole-cell pertussis (eg, DTP) or an acellular pertussis component, the decision to give ADACEL vaccine should be based on careful consideration of the potential benefits and possible risks: (2) (3)

- Temperature of 40.5°C (105°F) within 48 hours not due to another identifiable cause;
- Collapse or shock-like state (hypotonic-hyporesponsive episode) within 48 hours;
- Persistent, inconsolable crying lasting 3 hours, occurring within 48 hours;
- Seizures with or without fever occurring within 3 days.

When a decision is made to withhold pertussis vaccine, Td vaccine should be given. Persons who experienced Arthus-type hypersensitivity reactions (eg, severe local reactions associated with systemic symptoms) (4) following a prior dose of tetanus toxoid usually have high serum tetanus antitoxin levels and should not be given emergency doses of tetanus toxoid-containing vaccines more frequently than every 10 years, even if the wound is neither clean nor minor. (4) (5) If Guillain-Barré Syndrome occurred within 6 weeks of receipt of prior vaccine containing tetanus toxoid, the decision to give ADACEL vaccine or any vaccine containing tetanus toxoid should be based on careful consideration of the potential benefits and possible risks. (1) The decision to administer a pertussis-containing vaccine to individuals with stable central nervous system (CNS) disorders must be made by the health-care provider on an individual basis, with consideration of all relevant factors and assessment of potential risks and benefits for that individual. The ACIP has issued guidelines for immunizing such individuals. (2) A family history of seizures or other CNS disorders is not a contraindication to pertussis vaccine. (2) The ACIP has published guidelines for vaccination of persons with recent or acute illness. (1)

PRECAUTIONS General Do not administer by intravascular injection; ensure that the needle does not penetrate a blood vessel. ADACEL vaccine should not be administered into the buttocks nor by the intradermal route, since these methods of administration have not been studied; a weaker immune response has been observed when these routes of administration have been used with other vaccines. (1) The possibility of allergic reactions in persons sensitive to components of the vaccine should be evaluated. Epinephrine Hydrochloride Solution (1:1,000) and other appropriate agents and equipment should be available for immediate use in case of anaphylactic or acute hypersensitivity reaction occurs. Prior to administration of ADACEL vaccine, the vaccine recipient and/or the parent or guardian must be asked about personal health history, including immunization history, current health status and any adverse event after previous immunizations. In persons who have a history of serious or severe reaction within 48 hours of a previous injection with a vaccine containing similar components, administration of ADACEL vaccine must be carefully considered. The ACIP has published guidelines for the immunization of immunocompromised individuals. (6) Immune responses to inactivated vaccines and toxoids when given to immunocompromised persons may be suboptimal. (1) The immune response to ADACEL vaccine administered to immunocompromised persons (whether from disease or treatment) has not been studied. A separate, sterile syringe and needle, or a sterile disposable unit, must be used for each person to prevent transmission of blood borne infectious agents. Needles should not be recapped but should be disposed of according to biohazard waste guidelines.

Information for Vaccine Recipients and/or Parent or Guardian Before administration of ADACEL vaccine, health-care providers should inform the vaccine recipient and/or parent or guardian of the benefits and risks. The health-care provider should inform the vaccine recipient and/or parent or guardian about the potential for adverse reactions that have been temporally associated with ADACEL vaccine or other vaccines containing similar components. The vaccine recipient and/or parent or guardian should be instructed to report any serious adverse reactions to their health-care provider. Females of childbearing potential should be informed that Sanofi Pasteur Inc. maintains a pregnancy registry to monitor fetal outcomes of pregnant women exposed to ADACEL vaccine. If they are pregnant or become aware they were pregnant at the time of ADACEL vaccine immunization, they should contact their health-care professional or Sanofi Pasteur Inc. at 1-800-822-2463 (1-800-VACCINE). The health-care provider should provide the Vaccine Information Statements (VIS) that are required by the National Childhood Vaccine Injury Act of 1986 to be given with each immunization. The US Department of Health and Human Services has established a Vaccine Adverse Event Reporting System (VAERS) to accept all reports of suspected adverse events after the administration of any vaccine, including but not limited to the reporting of events required by the National Childhood Vaccine Injury Act of 1986. (7) The toll-free number for VAERS forms and information is 1-800-822-7967 or visit the VAERS website at <http://www.fda.gov/cber/vaers/vaers.htm>

Drug Interactions Immunosuppressive therapies, including irradiation, antimetabolites, alkylating agents, cytotoxic drugs and corticosteroids (used in greater than physiologic doses), may reduce the immune response to vaccines. (See PRECAUTIONS, General.) For information regarding simultaneous administration with other vaccines refer to the ADVERSE REACTIONS and DOSAGE AND ADMINISTRATION sections.

Carcinogenesis, Mutagenesis, Impairment of Fertility No studies have been performed with ADACEL vaccine to evaluate carcinogenicity, mutagenic potential, or impairment of fertility.

Pregnancy Category C Animal reproduction studies have not been conducted with ADACEL vaccine. It is also not known whether ADACEL vaccine can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. ADACEL vaccine should be given to a pregnant woman only if clearly needed. Animal fertility studies have not been conducted with ADACEL vaccine. The effect of ADACEL vaccine on embryo-fetal and pre-weaning development was evaluated in two developmental toxicity studies using pregnant rabbits. Animals were administered ADACEL vaccine twice prior to gestation, during the period of organogenesis (gestation day 6) and later during pregnancy on gestation day 29, 0.5 mL/rabbit/occasion (a 17-fold increase compared to the human dose of ADACEL vaccine on a body weight basis), by intramuscular injection. No adverse effects on pregnancy, parturition, lactation, embryo-fetal or pre-weaning development were observed. There were no vaccine related fetal malformations or other evidence of teratogenesis noted in this study. (8)

Pregnancy Registry Health-care providers are encouraged to register pregnant women who receive ADACEL vaccine in Sanofi Pasteur Inc.'s vaccination pregnancy registry by calling 1-800-822-2463 (1-800-VACCINE).

Nursing Mothers It is not known whether ADACEL vaccine is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when ADACEL vaccine is given to a nursing woman.

Pediatric Use ADACEL vaccine is not indicated for individuals less than 11 years of age. (See INDICATIONS AND USAGE.) For immunization of persons 6 weeks through 6 years of age against diphtheria, tetanus and pertussis refer to manufacturers' package inserts for DTap vaccines.

Geriatric Use ADACEL vaccine is not indicated for individuals 65 years of age and older. No data are available regarding the safety and effectiveness of ADACEL vaccine in individuals 65 years of age and older as clinical studies of ADACEL vaccine did not include subjects in the geriatric population.

ADVERSE REACTIONS The safety of ADACEL vaccine was evaluated in 4 clinical studies. A total of 5,841 individuals 11-64 years of age inclusive (3,393 adolescents 11-17 years of age and 2,448 adults 18-64 years) received a single booster dose of ADACEL vaccine. The principal safety study was a randomized, observer blind, active controlled trial that enrolled participants 11-17 years of age (ADACEL vaccine N = 1,184; Td vaccine N = 792) and 18-64 years of age (ADACEL vaccine N = 1,752; Td vaccine N = 573). Study participants had not received tetanus or diphtheria containing vaccines within the previous 5 years. Observer blind design, ie, study personnel collecting the safety data differed from personnel administering the vaccines, was used due to different vaccine packaging (ADA-

R only

CEL vaccine supplied in single dose vials; Td vaccine supplied in multi-dose vials). Solicited local and systemic reactions and unsolicited events were monitored daily for 14 days post-vaccination using a diary card. From days 14-28 post-vaccination, information on adverse events necessitating a medical contact, such as a telephone call, visit to an emergency room, physician's office or hospitalization, was obtained via telephone interview or at an interim clinic visit. From days 28 to 6 months post-vaccination, participants were monitored for unexpected visits to a physician's office or to an emergency room, onset of serious illness and hospitalizations. Information regarding adverse events that occurred in the 6 month post-vaccination time period was obtained via a scripted telephone interview. Approximately 96% of participants completed the 6-month follow-up evaluation. In the concomitant vaccination study with ADACEL and Hepatitis B vaccines, local and systemic adverse events were monitored daily for 14 days post-vaccination using a diary card. Local adverse events were only monitored at site/arm of ADACEL vaccine administration. Unsolicited reactions (including immediate reactions, serious adverse events and events that elicited seeking medical attention) were collected at a clinic visit or via telephone interview for the duration of the trial, ie, up to six months post-vaccination. In the concomitant vaccination study with ADACEL vaccine and trivalent inactivated influenza vaccines local and systemic adverse events were monitored for 14 days post-vaccination using a diary card. All unsolicited reactions occurring through day 14 were collected. From day 14 to the end of the trial, ie, up to 84 days, only events that elicited seeking medical attention were collected. In all studies, subjects were monitored for serious adverse events throughout the duration of the study. Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a vaccine cannot be directly compared to rates in the clinical trials of another vaccine and may not reflect the rates observed in practice. The adverse reaction information from clinical trials does, however, provide a basis for identifying the adverse events that appear to be related to vaccine use and for approximating rates of those events.

Serious Adverse Events in All Safety Studies Throughout the 6-month follow-up period in the principal safety study, serious adverse events were reported in 1.5% of ADACEL vaccine recipients and 1.4% in Td vaccine recipients. Two serious adverse events in adults were neurological events that occurred within 28 days of ADACEL vaccine administration; one serious migraine with unilateral facial paralysis and one diagnosis of nerve compression in neck and left arm. Similar or lower rates of serious adverse events were reported in the other trials and there were no additional neurological events reported.

Solicited Adverse Events in the Principal Safety Study The frequency of selected solicited adverse events (erythema, swelling, pain and fever) occurring during Days 0-14 following one dose of ADACEL vaccine or Td vaccine were reported at a similar frequency in both groups. Few participants (<1%) sought medical attention for these reactions. Pain at the injection site was the most common adverse reaction occurring in 62-78% of all vaccinees. In addition, overall rates of pain were higher in adolescent recipients of ADACEL vaccine compared to Td vaccine recipients. Rates of moderate and severe pain in adolescents did not significantly differ between the two groups. Rates of pain did not significantly differ for adults. Fever of 38°C and higher was uncommon, although in the adolescent age group, it occurred significantly more frequently in ADACEL vaccine recipients than Td vaccine recipients. (8) The rates of other local and systemic solicited reactions occurred at similar rates in ADACEL vaccine and Td vaccine recipients in the 3 day post-vaccination period. Most local reactions occurred within the first 3 days after vaccination (with a mean duration of less than 3 days). Headache was the most frequent systemic reaction and was usually of mild to moderate intensity.

Adverse Events in the Concomitant Vaccine Studies

Local and Systemic Reactions when Given with Hepatitis B Vaccine The rates reported for fever and injection site pain (at the ADACEL vaccine administration site) were similar when ADACEL and Hep B vaccines were given concurrently or separately. However, the rates of injection site erythema (23.4% for concomitant vaccination and 21.4% for separate administration) and swelling (23.9% for concomitant vaccination and 17.9% for separate administration) at the ADACEL vaccine administration site were increased when co-administered. Swollen and/or sore joints were reported by 22.5% for concomitant vaccination and 17.9% for separate administration. The rates of generalized body aches in the individuals who reported swollen and/or sore joints were 86.7% for concomitant vaccination and 72.2% for separate administration. Most joint complaints were mild in intensity with a mean duration of 1.8 days. The incidence of other solicited and unsolicited adverse events were not different between the 2 study groups. (8)

Local and Systemic Reactions when Given with Trivalent Inactivated Influenza Vaccine The rates of fever and injection site erythema and swelling were similar for recipients of concurrent and separate administration of ADACEL vaccine and TIV. However, pain at the ADACEL vaccine injection site occurred at statistically higher rates following concurrent administration (66.6%) versus separate administration (60.8%). The rates of sore and/or swollen joints were 13% for concurrent administration and 9% for separate administration. Most joint complaints were mild in intensity with a mean duration of 2.0 days. The incidence of other solicited and unsolicited adverse events were similar between the 2 study groups. (8)

Additional Studies An additional 1,806 adolescents received ADACEL vaccine as part of the lot consistency study used to support ADACEL vaccine licensure. This study was a randomized, double-blind, multi-center trial designed to assess lot consistency as measured by the safety and immunogenicity of 3 lots of ADACEL vaccine when given as a booster dose to adolescents 11-17 years of age inclusive. Local and systemic adverse events were monitored for 14 days post-vaccination using a diary card. Unsolicited adverse events and serious adverse events were collected for 28 days post-vaccination. Pain was the most frequently reported local adverse event occurring in approximately 80% of all subjects. Headache was the most frequently reported systemic event occurring in approximately 44% of all subjects. Sore and/or swollen joints were reported by approximately 14% of participants. Most joint complaints were mild in intensity with a mean duration of 2.0 days. (8) An additional 962 adolescents and adults received ADACEL vaccine in three supportive Canadian studies used as the basis for licensure in other countries. Within these clinical trials, the rates of local and systemic reactions following ADACEL vaccine were similar to those reported in the four principal trials in the US with the exception of a higher rate (86%) of adults experiencing 'any' local injection site pain. The rate of severe pain (0.8%), however, was comparable to the rates reported in the four principal trials. (8) There was one spontaneous report of whole-arm swelling of the injected limb among the 277 Td vaccine recipients, and two spontaneous reports among the 962 ADACEL vaccine recipients.

Postmarketing Reports The following adverse events have been spontaneously reported during the post-marketing use of ADACEL vaccine in other countries. Because these events are reported voluntarily from a population of uncertain size, it is not possible to reliably estimate their frequency or establish a causal relationship to vaccine exposure. The following adverse events were included based on severity, frequency of reporting or the strength of causal association to ADACEL vaccine. General disorders and administration site conditions: injection site bruising, sterile abscess; skin and subcutaneous tissue disorders: pruritus, urticaria. There have been spontaneous reports of nervous system disorders such as myelitis, syncope vasovagal, paresthesia, hyposthesia and musculoskeletal and connective tissue disorders such as myositis and muscle spasms temporally associated with ADACEL vaccine.

Reporting of Adverse Events The National Vaccine Injury Compensation Program, established by the National Childhood Vaccine Injury Act of 1986, requires physicians and other health-care providers who administer vaccines to maintain permanent vaccination records of the manufacturer and lot number of the vaccine administered in the vaccine recipient's permanent medical record along with the date of administration of the vaccine and the name, address and title of the person administering the vaccine. The Act further requires the health-care professional to report to the US Department of Health and Human Services the occurrence following immunization of any event set forth in the Vaccine Injury Table. These include anaphylaxis or anaphylactic shock within 7 days; brachial neuritis within 28 days; an acute complication or sequelae (including death) of an illness, disability, injury, or condition referred to above, or any events that would contraindicate further doses of vaccine, according to this ADACEL vaccine package insert. (7) (9) (10) The US Department of Health and Human Services has established the Vaccine Adverse Event Reporting System (VAERS) to accept all reports of suspected adverse events after the administration of any vaccine. Reporting of all adverse events occurring after vaccine administration is encouraged from vaccine recipients, parents/guardians and the health-care provider. Adverse events following immunization should be reported to VAERS. Reporting forms and information about reporting requirements or completion of the form can be obtained from VAERS through a toll-free number 1-800-822-7967 or visit the VAERS website at <http://www.fda.gov/cber/vaers/vaers.htm>. (7) (9) (10) Health-care providers should also report these events to Pharmacovigilance Department, Sanofi Pasteur Inc., Discovery Drive, Swiftwater, PA 18370 or call 1-800-822-2463 (1-800-VACCINE).

DOSAGE AND ADMINISTRATION ADACEL vaccine should be administered as a single injection of one dose (0.5 mL) by the intramuscular route. SHAKE THE VIAL WELL to distribute the suspension uniformly before withdrawing the 0.5 mL dose for administration. Five years should have elapsed since the recipient's last dose of tetanus toxoid, diphtheria toxoid and/or pertussis containing vaccine. Do NOT administer this product intravenously or subcutaneously.

STORAGE Store at 2° to 8°C (35° - 46°F). DO NOT FREEZE. Discard product if exposed to freezing. Do not use after expiration date.

REFERENCES 1. CDC. General recommendations on immunization: recommendations of the Advisory Committee on Immunization Practices (ACIP) and the American Academy of Family Physicians (AAFP). MMWR 2002;51(RR-2):1-35. 2. CDC. Pertussis vaccination: Use of acellular pertussis vaccines among infants and young children. Recommendations of the ACIP. MMWR 1997;46(RR-7):1-25. 3. CDC Update. Vaccine side effects, adverse reactions, contraindications and precautions - recommendations of the Advisory Committee on Immunization Practices (ACIP). MMWR 1996;45(RR-12):1-35. 4. CDC. Update on adult immunization: recommendations of the Advisory Committee on Immunization Practices (ACIP). MMWR 1991;40(RR-12):1-52. 5. CDC. Diphtheria, tetanus and pertussis: recommendations for vaccine use and other preventive measures. Recommendations of the Immunization Practices Advisory Committee (ACIP). MMWR 1991;40(RR-10):1-28. 6. CDC. Use of vaccines and immune globulins in persons with altered immunocompetence. Recommendations of the Advisory Committee on Immunization Practices (ACIP). MMWR 1993;42(RR-4):1-18. 7. CDC. Current trends - Vaccine Adverse Event Reporting System (VAERS) United States. MMWR 1990;39(41):730-3. 8. Data on file at Sanofi Pasteur Limited. 9. CDC. Current trends - national vaccine injury act: requirements for permanent vaccination records and for reporting of selected events after vaccination. MMWR 1988;37(13):197-200. 10. FDA. New reporting requirements for vaccine adverse events. FDA Drug Bull 1988;18(2):16-8.

Product information as of January 2006

Manufactured by:
Sanofi Pasteur Limited
Toronto Ontario Canada
MKT10383-1R

Printed in Canada

Distributed by:
Sanofi Pasteur Inc.
Swiftwater PA 18370 USA
R1-0106