

# Too Many Pregnant Women Still Drink Alcohol

BY MICHELE G. SULLIVAN

**P**regnant women consumed just as much alcohol in 2005 as they did in 1991, with 12% drinking at least once during pregnancy and 2% reporting binge drinking.

The findings illustrate the small effect of national educational programs aimed at decreasing this dangerous behavior, according to primary author Clark Denny, Ph.D. (MMWR 2009;58:529-32).

In Healthy People 2010, the national health agenda published in 2000, Dr. David Satcher, the U.S. surgeon general at that time, set abstinence targets of 95% for alcohol and 100% for binge drinking among pregnant women.

"The prevalence of both types of drinking behavior among pregnant women remains higher than the Healthy People 2010 targets and greater progress will be needed to reach them," Dr. Denny wrote.

The 15-year study found that women aged 35-44 years had the highest incidence of drinking during pregnancy (18%), wrote Dr. Denny, an epidemiologist from the Centers for Disease Control and Prevention. Rates were also higher in college-educated women, employed women, and unmarried women.

The study was based on data collected from 1991 to 2005 through the Behavioral Risk Factor Surveillance System surveys. These annual surveys randomly poll community-dwelling U.S. adults about behavioral health issues. The CDC study included data collected from women aged 18-44 years, who were asked about alcohol use (at least one drink in the last 30 days) and binge drinking (at least five drinks on any one occasion in the last 30 days).

During the 15-year period, 533,500 women were surveyed; 22,000 (4%) reported being pregnant at the time of the survey. The average annual percentage of any alcohol use among the pregnant women was 12%, and did not change from 1991 to 2005. The average annual percentage of pregnant women who said they binged was 2%; again, that percentage was stable over the survey period.

From 2001 to 2005, the study also examined the relationship between drinking during pregnancy and demographic factors. Age was associated with both any drinking and binge drinking. The oldest women (35-44 years) had the highest drinking rate (18%), while the youngest women (18-24 years) had the lowest rate (9%). Age was not highly associated with binge drinking.

Education, employment, and marital status were also associated with drinking during pregnancy. Any drinking was higher among employed women than unemployed (14% vs. 8%), and among unmarried women than married women (13% vs. 10%). Binge drinking was also more common among employed than unemployed women (2% vs. 1%), and unmarried women than married women (4% vs. 1%).

The reasons for these associations are

unclear, wrote Dr. Denny. "Some possible reasons include that older women might be more likely to be alcohol dependent and have more difficulty abstaining while pregnant, more educated and employed women might have more discretionary money for the purchase of alcohol, and unmarried women might attend more social occasions where alcohol is served."

Rates of drinking and binge drinking


were higher among nonpregnant women (54% and 12%). Prepregnancy alcohol use is a strong predictor of use during pregnancy, and many women who drink continue to do so before realizing that they are pregnant, Dr. Denny noted. "Approximately 40% of women realize they are pregnant at 4 weeks of gestation, a critical period for fetal organ development."

The findings confirm the need for al-

cohol use screening and counseling among all women, he wrote in a press statement. "By screening and advising women about the risks of drinking while pregnant, health care providers can play a key role in reducing rates of fetal alcohol syndrome. This study revealed that there is still a great need for health care professionals to routinely ask all women who are pregnant or at risk of being pregnant about their alcohol consumption." ■

## Why would the most trusted lubricant brand create an arousal gel?

**Because K-Y® Brand never stops pioneering ways to fulfill women's unmet sexual needs.**



Introducing K-Y® Brand INTENSE™: a female arousal gel developed in response to studies showing that more exhilarating orgasms top women's lists of unmet sexual needs. Three years of extensive research led K-Y® Brand scientists to develop a proprietary blend of niacin and sensory ingredients that have been clinically shown to heighten female sexual arousal and satisfaction when massaged onto the clitoris. In fact, 75% of women using K-Y® Brand INTENSE™ in consumer studies reported heightened sensitivity, arousal, and sexual pleasure. So for women looking for ways to enhance their sexual well-being, recommend K-Y® Brand INTENSE™: the first and only female arousal gel from K-Y® Brand, THE TRUSTED INTIMACY EXPERT™.

And for their other intimate needs, remind your patients that K-Y® Brand INTENSE™ is not a lubricant, but it can be combined with other K-Y® Brand lubricants and vaginal moisturizers, such as K-Y® Brand LIQUIBEADS™ with exclusive OVULE® technology or K-Y® Brand SENSITIVE™ Jelly.

**K-Y®** Applying science to  
the art of intimacy.  
BRAND • MARQUE

©McNEIL-PPC, Inc. 2009