

# No Lasting Effects From Repeated Ultrasounds

BY ROBERT FINN  
San Francisco Bureau

An 8-year follow-up study from a randomized, controlled trial of repeated prenatal ultrasound examinations found no evidence of any lasting harm to any aspect of child development.

The initial study demonstrated that infants exposed prenatally to five ultrasound imaging studies between 18 and 38 weeks' gestation had a significantly greater risk of low birth weight than children exposed to only a single ultrasound at 18 weeks of gestation.

Included in the follow-up analysis were 1,352 children who previously had been randomized to the regular ultrasound group and 1,362 children randomized to

the intensive ultrasound group (Lancet 2004;364:2038-44).

After 8 years of repeated follow-up examinations, children in the two groups showed no statistically significant differences in a wide variety of developmental measures. These included measures of physical growth, toddler temperament, language development, and behavior, reported John P. Newham, M.D., of the University of Western Australia (Subiaco), and his colleagues.

The groups showed a statistically significant difference on only a single measure of child development. At 1 year of age children in the intensive ultrasound group showed a smaller number of abnormal scores on a test of early language milestones than children in the regular ultrasound groups.

The authors suggested that this may have been a statistical fluke—a seemingly significant result that showed up by chance because of the many end points examined

in the study. An alternative explanation may be that women who had repeated ultrasound examinations may have had greater awareness of the study, which in turn enhanced parental attention, resulting in earlier language acquisition.

While this study showed no deleterious effects of repeated ultrasound examinations to the developing fetus, the authors cautioned that contemporary ultrasound instruments have higher power outputs than the instruments used in the study. ■

## Raised Serum Bile Acid Level Linked to IHP

VIENNA — An elevated serum bile acid level is a highly reliable indicator of intrahepatic cholestasis of pregnancy in a woman who presents with itching and excoriated skin lesions late in gestation, Christina M. Rudolph, M.D., reported at the annual meeting of the European Society for Dermatological Research.

**The lowest serum bile acid level in the ICP patients was markedly greater than the highest value in other pregnant women with pruritic conditions.**

The differential diagnosis of pruritic skin conditions during pregnancy has often been vexing because of extensive overlap in clinical presentation.

But in her series of 75 patients who presented with pruritic skin changes to a specialized dermatology clinic for pregnant women, the lowest serum bile acid level among the 11 patients with intrahepatic cholestasis of pregnancy (ICP)—7.3  $\mu\text{mol/L}$ —was markedly greater than the highest value among women with other pruritic conditions, said Dr. Rudolph of the University of Graz, Austria.


The distinction is clinically important because ICP, if untreated, is associated with increased risks of stillbirth and preterm delivery, she noted.

Other dermatologic diagnoses made in this cohort were atopy-related skin changes, specific dermatoses of pregnancy, psoriasis, pityriasis rosea, and drug reaction. The mean serum bile acid level in these women was 2.3  $\mu\text{mol/L}$ , with a range of 0.4-4.5  $\mu\text{mol/L}$ .

In contrast, the range of serum bile acid levels in women with ICP was 7.3-138  $\mu\text{mol/L}$ , with a mean value of 37.4  $\mu\text{mol/L}$ .

—Bruce Jancin

For the local treatment of atrophic vaginitis...



Premarin Vaginal Cream

## CREAM VS CLEAN

**VAGIFEM—a neat alternative to Premarin® Vaginal Cream (PVC)**

- Clean, leak-free estrogen delivery
- Symptom relief comparable to PVC\*1
- Greater patient acceptance versus PVC<sup>1</sup>
- Lower systemic absorption versus PVC<sup>1</sup>
- Generally mild adverse events

— Those most commonly reported ( $\geq 5\%$ ) include headache, abdominal pain, back pain, upper respiratory tract infection, and genital moniliasis

VAGIFEM should not be used in women with known or suspected breast cancer or estrogen-dependent neoplasia, abnormal genital bleeding of unknown etiology, known or suspected pregnancy, porphyria, hypersensitivity to any of its components, active or a history of thrombophlebitis or thromboembolic disorders, or a history of thrombosis.

ESTROGENS HAVE BEEN REPORTED TO INCREASE THE RISK OF ENDOMETRIAL CARCINOMA.

Premarin (conjugated estrogens) Vaginal Cream is a registered trademark of Wyeth Pharmaceuticals.

\*A 24-week, multicenter, randomized, open-label, controlled study of 159 postmenopausal women treated with VAGIFEM (n=80; 1 tablet/day for 2 weeks, then twice weekly for 22 weeks) or PVC (n=79; 2 g/day for 3 weeks, 1 week off, then repeated 3 weeks on and 1 week off for the rest of the study). Based on composite scores of 3 vaginal symptoms: dryness, soreness, and irritation.

Please see adjacent page for brief summary of Prescribing Information.

[www.vagifem.com](http://www.vagifem.com)



novo nordisk®

**VAGIFEM**<sup>25µg</sup>  
estradiol vaginal tablets  
NEATLY EFFECTIVE

