

Aspirin and Esomeprazole Appear Safe for Barrett's

BY FRAN LOWRY
Orlando Bureau

ORLANDO — Early findings from the Aspirin Esomeprazole Chemoprevention Trial indicate that therapy with aspirin and esomeprazole is safe and well tolerated for preventing the progression of Barrett's esophagus to adenocarcinoma.

Since the start of the randomized Aspirin Esomeprazole Chemoprevention Trial (AspECT) in September 2005, 1,192

(83%) of the 1,436 patients have remained on their medication, and just 33 adverse events have been reported, said the study's lead investigator Dr. Janusz Jankowski, professor of medicine, Oxford University (England), at a meeting on gastrointestinal cancers sponsored by the American Society of Clinical Oncology.

AspECT is an ambitious, 10-year clinical trial being conducted in the United Kingdom. The investigators are still recruiting to meet their goal of 3,000 pa-

tients. The trial's primary aim is to determine whether treatment with the proton pump inhibitor esomeprazole (Nexium, AstraZeneca) and aspirin can stop Barrett's metaplasia from progressing to adenocarcinoma.

The investigators are also trying to determine whether this therapy will prevent or reduce myocardial infarction.

The United Kingdom is fertile ground for such a study, Dr. Jankowski said at the symposium, also sponsored by the AGA

Institute, the American Society for Therapeutic Radiology and Oncology, and the Society of Surgical Oncology.

"The U.K. has the highest incidence of esophageal adenocarcinoma in the world—up to four times greater than that of other countries in Europe. Barrett's metaplasia is twice as common in the U.K. as it is in the United States," he said in an interview.

Being able to show that aspirin "is in-
Continued on following page

Tetanus Toxoid, Reduced Diphtheria Toxoid and Acellular Pertussis Vaccine Adsorbed

Adacel

Brief Summary: Please see package insert for full prescribing information

INDICATIONS AND USAGE ADACEL™ vaccine is indicated for active booster immunization for the prevention of tetanus, diphtheria and pertussis as a single dose in persons 11 through 64 years of age. The use of ADACEL vaccine as a primary series, or to complete the primary series, has not been studied. As with any vaccine, ADACEL vaccine may not protect 100% of vaccinated individuals.

CONTRAINDICATIONS Known systemic hypersensitivity to any component of ADACEL vaccine or a life-threatening reaction after previous administration of the vaccine or a vaccine containing the same substances are contraindications to vaccination with ADACEL vaccine. Because of uncertainty as to which component of the vaccine may be responsible, additional vaccinations with the diphtheria, tetanus or pertussis components should not be administered. Alternatively, such individuals may be referred to an allergist for evaluation if further immunizations are to be considered. The following events are contraindications to administration of any pertussis containing vaccine: (1)

- Encephalopathy within 7 days of a previous dose of pertussis containing vaccine not attributable to another identifiable cause.
- Progressive neurological disorder, uncontrolled epilepsy, or progressive encephalopathy. Pertussis vaccine should not be administered to individuals with these conditions until a treatment regimen has been established, the condition has stabilized, and the benefit clearly outweighs the risk.

ADACEL vaccine is not contraindicated for use in individuals with HIV infection. (1)

WARNINGS Because intramuscular injection can cause injection site hematoma, ADACEL vaccine should not be given to persons with any bleeding disorder, such as hemophilia or thrombocytopenia, or to persons on anticoagulant therapy unless the potential benefits clearly outweigh the risk of administration. If the decision is made to administer ADACEL vaccine in such persons, it should be given with caution, with steps taken to avoid the risk of hematoma formation following injection. (1) If any of the following events occurred in temporal relation to previous receipt of a vaccine containing a whole-cell pertussis (eg, DTP) or an acellular pertussis component, the decision to give ADACEL vaccine should be based on careful consideration of the potential benefits and possible risks: (2)(3)

- Temperature of $\geq 40.5^{\circ}\text{C}$ (105°F) within 48 hours not due to another identifiable cause;
- Collapse or shock-like state (hypotonic-hyporesponsive episode) within 48 hours;
- Persistent, inconsolable crying lasting ≥ 3 hours, occurring within 48 hours;
- Seizures with or without fever occurring within 3 days.

When a decision is made to withhold pertussis vaccine, Td vaccine should be given. Persons who experienced Arthus-type hypersensitivity reactions (eg, severe local reactions associated with systemic symptoms) (4) following a prior dose of tetanus toxoid usually have high serum tetanus antitoxin levels and should not be given emergency doses of tetanus toxoid-containing vaccines more frequently than every 10 years, even if the wound is neither deep nor minor. (4)(5) If Guillain-Barré syndrome occurred within 6 weeks of receipt of prior vaccine containing tetanus toxoid, the decision to give ADACEL vaccine or any vaccine containing tetanus toxoid should be based on careful consideration of the potential benefits and possible risks. (1) The decision to administer a pertussis-containing vaccine to individuals with stable central nervous system (CNS) disorders must be made by the health-care provider on an individual basis, with consideration of all relevant factors and assessment of potential risks and benefits for that individual. The Advisory Committee on Immunization Practices (ACIP) has issued guidelines for immunizing such individuals. (2) A family history of seizures or other CNS disorders is not a contraindication to pertussis vaccine. (2) The ACIP has published guidelines for vaccination of persons with recent or acute illness. (1)

PRECAUTIONS **General** Do not administer by intravascular injection; ensure that the needle does not penetrate a blood vessel. ADACEL vaccine should not be administered into the buttocks or by the intrathecal route, since these methods of administration have not been studied; a weaker immune response has been observed when these routes of administration have been used with other vaccines. (1) The possibility of allergic reactions in persons sensitive to components of the vaccine should be evaluated. Epinephrine Hydrochloride Solution (1:1,000) and other appropriate agents and equipment should be available for immediate use in case an anaphylactic or acute hypersensitivity reaction occurs. Prior to administration of ADACEL vaccine, the vaccine recipient and/or the parent or guardian must be asked about personal health history, including immunization history, current health status and any adverse event after previous immunizations. In persons who have a history of serious or severe reaction within 48 hours of a previous injection with a vaccine containing similar components, administration of ADACEL vaccine must be carefully considered. The ACIP has published guidelines for the immunization of immunocompromised individuals. (6) Immune responses to inactivated vaccines and toxoids when given to immunocompromised persons may be suboptimal. (1) The immune response to ADACEL vaccine administered to immunocompromised persons (whether from disease or treatment) has not been studied. A separate, sterile syringe and needle, or a sterile disposable unit, must be used for each person to prevent transmission of blood borne infectious agents. Needles should not be re-sterilized but should be disposed of according to biohazard waste guidelines.

Information for Vaccine Recipients and/or Parent or Guardian Before administration of ADACEL vaccine, health-care providers should inform the vaccine recipient and/or parent or guardian of the benefits and risks. The health-care provider should inform the vaccine recipient and/or parent or guardian about the potential for adverse reactions that have been temporally associated with ADACEL vaccine or other vaccines containing similar components. The vaccine recipient and/or parent or guardian should be instructed to report any serious adverse reactions to their health-care provider. Females of childbearing potential should be informed that Sanofi Pasteur Inc. maintains a pregnancy registry to monitor fetal outcomes of pregnant women exposed to ADACEL vaccine. If they are pregnant or become aware they were pregnant at the time of ADACEL vaccine immunization, they should contact their health-care professional or Sanofi Pasteur Inc. at 1-800-822-2463 (1-800-VACCINE). The health-care provider should provide the Vaccine Information Statements (VIS) that are required by the National Childhood Vaccine Injury Act of 1986 to be given with each immunization. The US Department of Health and Human Services has established a Vaccine Adverse Event Reporting System (VAERS) to accept all reports of suspected adverse events after the administration of any vaccine, including but not limited to the reporting of events required by the National Childhood Vaccine Injury Act of 1986. (7) The toll-free number for VAERS forms and information is 1-800-822-7967 or visit the VAERS website at <http://www.fda.gov/cber/vaers/vaers.htm>

Drug Interactions Immunosuppressive therapies, including irradiation, antimetabolites, alkylating agents, cytotoxic drugs and corticosteroids (used in greater than physiologic doses), may reduce the immune response to vaccines. (See **PRECAUTIONS, General**.) For information regarding simultaneous administration with other vaccines refer to the **ADVERSE REACTIONS AND DOSAGE AND ADMINISTRATION** sections.

Carcinogenesis, Mutagenesis, Impairment of Fertility No studies have been performed with ADACEL vaccine to evaluate carcinogenicity, mutagenic potential, or impairment of fertility.

Pregnancy Category C Animal reproduction studies have not been conducted with ADACEL vaccine. It is also not known whether ADACEL vaccine can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. ADACEL vaccine should be given to a pregnant woman only if clearly needed. Animal fertility studies have not been conducted with ADACEL vaccine. The effect of ADACEL vaccine on embryo-fetal and pre-weaning development was evaluated in two developmental toxicity studies using pregnant rabbits. Animals were administered ADACEL vaccine twice prior to gestation, during the period of organogenesis (gestation day 6) and later during pregnancy on gestation day 29. 0.5 mL/rabbit/occasion (a 17-fold increase compared to the human dose of ADACEL vaccine on a body weight basis), by intramuscular injection. No adverse effects on pregnancy, parturition, lactation, embryo-fetal or pre-weaning development were observed. There were no vaccine related fetal malformations or other evidence of teratogenic noted in this study. (8)

Pregnancy Registry Health-care providers are encouraged to register pregnant women who receive ADACEL vaccine in Sanofi Pasteur Inc.'s vaccination pregnancy registry by calling 1-800-822-2463 (1-800-VACCINE).

Nursing Mothers It is not known whether ADACEL vaccine is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when ADACEL vaccine is given to a nursing woman.

Pediatric Use ADACEL vaccine is not indicated for individuals less than 11 years of age. (See **INDICATIONS AND USAGE**.) For immunization of persons 6 weeks through 6 years of age against diphtheria, tetanus and pertussis refer to manufacturers' package inserts for DTPa vaccines.

Geriatric Use ADACEL vaccine is not indicated for individuals 65 years of age and older. No data are available regarding the safety and effectiveness of ADACEL vaccine in individuals 65 years of age and older as clinical studies of ADACEL vaccine did not include subjects in the geriatric population.

ADVERSE REACTIONS The safety of ADACEL vaccine was evaluated in 4 clinical studies. A total of 5,841 individuals 11-64 years of age inclusive (3,393 adolescents 11-17 years of age and 2,448 adults 18-64 years) received a single booster dose of ADACEL vaccine. The principal safety study was a randomized, observer blind, active controlled trial that enrolled participants 11-17 years of age (ADACEL vaccine N = 1,184; Td vaccine N = 792) and 18-64 years of age (ADACEL vaccine N = 1,752; Td vaccine N = 573). Study participants had not received tetanus or diphtheria containing vaccines within the previous 5 years. Observer blind design, ie, study

Product information as of January 2006

Manufactured by:
Sanofi Pasteur Limited
Toronto Ontario Canada
MKT14427

personnel collecting the safety data differed from personnel administering the vaccines, was used due to different vaccine packaging (ADACEL vaccine supplied in single dose vials; Td vaccine supplied in multi-dose vials). Solicited local and systemic reactions and unsolicited events were monitored daily for 14 days post-vaccination using a diary card. From days 14-28 post-vaccination, information on adverse events necessitating a medical contact, such as a telephone call, visit to an emergency room, physician's office or hospitalization, was obtained via telephone interview or at an interim clinic visit. From days 28 to 6 months post-vaccination, participants were monitored for unexpected visits to a physician's office or to an emergency room, onset of serious illness and hospitalizations. Information regarding adverse events that occurred in the 6 month post-vaccination time period was obtained via a scripted telephone interview. Approximately 96% of participants completed the 6 month follow-up evaluation. In the concomitant vaccination study with ADACEL and Hepatitis B vaccines, local and systemic adverse events were monitored daily for 14 days post-vaccination using a diary card. Local adverse events were only monitored at site/arm of ADACEL vaccine administration. Unsolicited reactions (including immediate reactions, serious adverse events and events that elicited seeking medical attention) were collected at a clinic visit or via telephone interview for the duration of the trial, ie, up to 6 months post-vaccination. In the concomitant vaccination study with ADACEL vaccine and trivalent inactivated influenza vaccines, local and systemic adverse events were monitored for 14 days post-vaccination using a diary card. All unsolicited reactions occurring through day 14 were collected. From day 14 to the end of the trial, ie, up to 84 days, only events that elicited seeking medical attention were collected. In all the studies, subjects were monitored for serious adverse events throughout the duration of the study. Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a vaccine cannot be directly compared to rates in the clinical trials of another vaccine and may not reflect the rates observed in practice. The adverse reaction information from clinical trials does, however, provide a basis for identifying the adverse events that appear to be related to vaccine use and for approximating rates of those events.

Serious Adverse Events in All Safety Studies Throughout the 6-month follow-up period in the principal safety study, serious adverse events were reported in 1.5% of ADACEL vaccine recipients and 1.4% in Td vaccine recipients. Two serious adverse events in adults were neurologic events that occurred within 28 days of ADACEL vaccine administration: one severe migraine with unilateral facial paralysis and one diagnosis of nerve compression in neck and left arm. Similar or lower rates of serious adverse events were reported in the other trials and there were no additional neurologic events reported.

Solicited Adverse Events in the Principal Safety Study The frequency of selected solicited adverse events (erythema, swelling, pain and fever) occurring during Days 0-14 following one dose of ADACEL vaccine or Td vaccine were reported at a similar frequency in both groups. Few participants (<1%) sought medical attention for these reactions. Pain at the injection site was the most common adverse reaction occurring in 62-78% of all vaccinees. In addition, overall rates of pain were higher in adolescent recipients of ADACEL vaccine compared to Td vaccine recipients. Rates of moderate and severe pain in adolescents did not significantly differ between the two groups. Rates of pain did not significantly differ for adults. Fever of 38°C and higher was uncommon, although in the adolescent age group, it occurred significantly more frequently in ADACEL vaccine recipients than Td vaccine recipients. (8) Headache was the most frequent systemic reaction and was usually of mild to moderate intensity. Local and systemic solicited reactions occurred at similar rates in ADACEL vaccine and Td vaccine recipients in the 3 day post-vaccination period. Most local reactions occurred within the first 3 days after vaccination (with a mean duration of less than 3 days).

Adverse Events in the Concomitant Vaccine Studies

Local and Systemic Reactions when Given with Hepatitis B Vaccine The rates reported for fever and injection site pain (at the ADACEL vaccine administration site) were similar when ADACEL and Hep B vaccines were given concurrently or separately. However, the rates of injection site erythema (23.4% for concomitant vaccination and 21.4% for separate administration) and swelling (23.9% for concomitant vaccination and 17.9% for separate administration) at the ADACEL vaccine administration site were increased when co-administered. Swollen and/or sore joints were reported by 22.5% for concomitant vaccination and 17.9% for separate administration. The rates of generalized body aches in the individuals who reported swollen and/or sore joints were 86.7% for concomitant vaccination and 72.2% for separate administration. Most joint complaints were mild in intensity with a mean duration of 1.8 days. The incidence of other solicited and unsolicited adverse events were not different between the 2 study groups. (8)

Local and Systemic Reactions when Given with Trivalent Inactivated Influenza Vaccine The rates of fever and injection site erythema and swelling were similar for recipients of concurrent and separate administration of ADACEL vaccine and Td vaccine. However, pain at the ADACEL vaccine injection site occurred at statistically higher rates following concurrent administration (65.6%) versus separate administration (60.8%). The rates of sore and/or swollen joints were 13% for concurrent administration and 9% for separate administration. Most joint complaints were mild in intensity with a mean duration of 2.0 days. The incidence of other solicited and unsolicited adverse events were similar between the 2 study groups. (8)

Additional Studies An additional 1,806 adolescents received ADACEL vaccine as part of the lot consistency study used to support ADACEL vaccine licensure. This study was a randomized, double-blind, multi-center trial designed to assess lot consistency as measured by the safety and immunogenicity of 3 lots of ADACEL vaccine when given as a booster dose to adolescents 11-17 years of age inclusive. Local and systemic adverse events were monitored for 14 days post-vaccination using a diary card. Unsolicited adverse events and serious adverse events were collected for 28 days post-vaccination. Pain was the most frequently reported local adverse event occurring in approximately 80% of all subjects. Headache was the most frequently reported systemic event occurring in approximately 44% of all subjects. Sore and/or swollen joints were reported by approximately 14% of participants. Most joint complaints were mild in intensity with a mean duration of 2.0 days. (8) An additional 962 adolescents and adults received ADACEL vaccine in three supportive Canadian studies used as the basis for licensure in other countries. Within these clinical trials, the rates of local and systemic reactions following ADACEL vaccine were similar to those reported in the four principal trials in the US with the exception of a higher rate (86%) of adults experiencing "any" local injection site pain. The rate of severe pain (0.8%), however, was comparable to the rates reported in the four principal trials. (8) There was one spontaneous report of whole-arm swelling of the injected limb among the 277 Td vaccine recipients, and two spontaneous reports among the 962 ADACEL vaccine recipients.

Postmarketing Reports The following adverse events have been spontaneously reported during the post-marketing use of ADACEL vaccine in other countries. Because these events are reported voluntarily from a population of uncertain size, it is not possible to reliably estimate their frequency or establish a causal relationship to vaccine exposure. The following adverse events were included based on severity, frequency of reporting or the strength of causal association to ADACEL vaccine. General disorders and administration site conditions: injection site bruise, sterile abscess. Skin and subcutaneous tissue disorders: pruritus, urticaria. There have been spontaneous reports of nervous system disorders such as myelitis, syncope, vasovagal, paresthesia, hypoesthesia and musculoskeletal and connective tissue disorders such as myositis and muscle spasms temporally associated with ADACEL vaccine.

Reporting of Adverse Events The National Vaccine Injury Compensation Program, established by the National Childhood Vaccine Injury Act of 1986, requires physicians and other health-care providers who administer vaccines to maintain permanent vaccination records of the manufacturer and lot number of the vaccine administered in the vaccine recipient's permanent medical record along with the date of administration of the vaccine and the name, address and title of the person administering the vaccine. The Act further requires the health-care professional to report to the US Department of Health and Human Services the occurrence following immunization of any event set forth in the Vaccine Injury Table. These include anaphylaxis or anaphylactic shock within 7 days; brachial neuritis within 28 days; an acute complication or sequelae (including death) of an illness, disability, injury, or condition related to above, or any events that would contraindicate further doses of vaccine, according to this ADACEL vaccine package insert. (7)(9)(10) The US Department of Health and Human Services has established the Vaccine Adverse Event Reporting System (VAERS) to accept all reports of suspected adverse events after the administration of any vaccine. Reporting of all adverse events occurring after vaccine administration is encouraged from vaccine recipients, parents/guardians and the health-care provider. Adverse events following immunization should be reported to VAERS. Reporting forms and information about reporting requirements or completion of the form can be obtained from VAERS through a toll-free number 1-800-822-7967 or visit the VAERS website at <http://www.fda.gov/cber/vaers/vaers.htm>. (7)(9)(10) Health-care providers should also report these events to the Pharmacovigilance Department, Sanofi Pasteur Inc., Discovery Drive, Swiftwater, PA 18370 or call 1-800-822-2463 (1-800-VACCINE).

DOSAGE AND ADMINISTRATION ADACEL vaccine should be administered as a single injection of one dose (0.5 mL) by the intramuscular route. SHAKE THE VIAL WELL to distribute the suspension uniformly before withdrawing the 0.5 mL dose for administration. Do NOT administer this product intravenously or subcutaneously. Five years should have elapsed since the recipient's last dose of tetanus toxoid, diphtheria toxoid and/or pertussis containing vaccine.

STORAGE Store at 2° to 8°C (35° to 46°F). DO NOT FREEZE. Discard product if exposed to freezing. Do not use after expiration date.

REFERENCES 1. Centers for Disease Control and Prevention (CDC). General recommendations on immunization: recommendations of the Advisory Committee on Immunization Practices (ACIP) and the American Academy of Family Physicians (AAFP). *MMWR* 2002;51(RR-2):1-35. 2. CDC. Pertussis vaccination: use of acellular pertussis vaccines among infants and young children. Recommendations of the ACIP. *MMWR* 1997;46(RR-7):1-25. 3. CDC Update. Vaccine side effects, adverse reactions, contraindications and precautions - recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR* 1996;45 (RR-12):1-35. 4. CDC. Update on adult immunization: recommendations of the ACIP. *MMWR* 1991;40(RR-12):1-52. 5. CDC. Diphtheria, tetanus and pertussis: recommendations for vaccine use and other preventive measures. Recommendations of the ACIP. *MMWR* 1991;40(RR-10):1-28. 6. CDC. Use of vaccines and immune globulins in persons with altered immunocompetence. Recommendations of the ACIP. *MMWR* 1993;42(RR-4):1-18. 7. CDC. Current trends - Vaccine Adverse Event Reporting System (VAERS) United States. *MMWR* 1990;39(41):730-3. 8. Data on file at Sanofi Pasteur Limited. 9. CDC. Current trends - national vaccine injury act: requirements for permanent vaccination records and for reporting of selected events after vaccination. *MMWR* 1988;37(13):197-200. 10. FDA. New reporting requirements for vaccine adverse events. *FDA Drug Bull* 1988;18(2):16-8.

Printed in Canada

Distributed by:
Sanofi Pasteur Inc.
Swiftwater PA 18370 USA
R2-0206USA
D82-372MQ
2023657-306

Tenofovir Beats Adefovir at Hep B Viral Suppression

BOSTON — Tenofovir suppresses viral load more rapidly and effectively than adefovir does in patients with HBe antigen-negative chronic hepatitis B, Dr. Patrick Marcellin reported at the annual meeting of the American Association for the Study of Liver Diseases.

Although both patient groups experienced a rapid decline in viral load by week 4 of the 48-week trial, those taking tenofovir experienced a steeper decline and a higher response rate, and the response was maintained, said Dr. Marcellin of the Hospital Beaujon, Clichy, France.

In the phase III trial, 375 patients with chronic hepatitis B infection were randomized to either 300 mg/day tenofovir or 10 mg/day adefovir. The primary end points were suppression of viral DNA to below 400 copies/mL and reduction of at least 2 points in the Knodell necroinflammatory score without worsening of fibrosis.

The patients' mean age was 44 years, and their mean necroinflammation score was 8. Twenty percent had cirrhosis. At baseline, mean hepatitis B virus RNA levels were about $7 \log_{10} \text{ c/mL}$.

Both groups achieved rapid suppression of hepatitis B virus DNA, with the majority of responsive patients doing so by week 4. By week 48, however, response differences emerged. Significantly more tenofovir-treated patients than adefovir-treated patients achieved viral loads below 400 copies/mL (93% versus 63%, respectively).

There was no significant difference in histologic response between the two groups: 72% treated with tenofovir improved, versus 69% treated with adefovir. But there was a significant difference in the percentage of patients who achieved both virologic and histologic response: 71% of the tenofovir group, versus 49% of the adefovir group.

At week 48, the ALT level was normal in 77% of both groups. The incidence of ALT flare was about 1% in each group. There were no significant differences in amylase, lipase, or creatinine levels. Regarding drug resistance, none of the tenofovir-treated patients developed resistant mutations.

The phase III study was sponsored by Gilead Sciences Inc., Durham, N.C., the company that manufactures tenofovir. Dr. Marcellin disclosed he has a financial relationship with Gilead Sciences.

—Michele G. Sullivan

Continued from previous page

credibly well tolerated" is very gratifying, Dr. Jankowski said, because many people were skeptical that it could be done.

"One of the major criticisms of the study when we tried to get it funded in the first place was that people thought we were mad and dangerous, and that we would kill patients with low-dose aspirin. But about 90% of our patients are still on low-dose aspirin and esomeprazole 2 years into the study with hardly any adverse events, showing the drug combination is very well tolerated."

So far, 12% of patients randomized to 20 mg esomeprazole have required an increase to 40 mg for symptom control, Dr. Jankowski said.

Besides conversion to high-grade dysplasia or cancer, the other primary end point of the study is all-cause mortality.

Additionally, it will look at the pharmacokinetics of aspirin resistance, genetic markers as potential risk factors for esophageal can-



cer, and quality of life.

The first planned efficacy analysis is scheduled for 2010, a second interim analysis is due in 2012, and the final analysis is due in 2016.

Being able to show that aspirin 'is incredibly well tolerated' is very gratifying because many people were skeptical.

DR. JANKOWSKI

a consultant to and receives research funding from AstraZeneca. ■

The trial is funded by Cancer Research UK, Oxford University, and AstraZeneca.

Dr. Jankowski disclosed that he is

Perception of Pain Altered in IBS Patients

BY KATE JOHNSON
Montreal Bureau

Patients with irritable bowel syndrome have altered brain responses to the anticipation of pain and to pain itself, which may make them more sensitive to painful stimuli.

"During expectation of pain, IBS [irritable bowel syndrome] patients generate higher levels of tonic noradrenergic activity, producing a bias toward interpretation of network activity as pain (speed over accuracy), and are inefficient at reducing such activity when discrimination of nonpainful stimulation should be maximized, or pain should be inhibited," wrote Dr. Steven M. Berman and his colleagues from the Center for the Neurobiology of Stress at the University of California, Los Angeles (J. Neurosci. 2008;28:349-59).

They used functional magnetic resonance imaging (fMRI) to measure the blood oxygen level-dependent response to anticipated and delivered rectal distention in 14 female IBS patients and 12 healthy controls (mean age 36 years). The imaging showed that, when the control subjects were anticipating a painful stimulus, brain activity decreased in the insula, supragenual anterior cingulate cortex, amygdala, and dorsal brainstem, but there was less of this anticipatory deactivation in the IBS patients.

Visceral distention of the rectum was then performed using a computer-driven pump and rectal balloon, which was inflated, in random order, to pressures of 25 mm Hg, 45 mm Hg, or 5 mm Hg (sham distention). The distention was performed after an 8-hour fast and two enemas.

Four to six sessions of 16 inflations were performed. Each inflation was preceded by an anticipatory cue, then 15 seconds of inflation at the designated pressure.

During rectal distention, increases in activity in the insula, dorsal anterior cingulate cortex, and dorsal brainstem were more extensive in IBS patients than in controls. "The DBS [dorsal brainstem] region contains multiple small structures implicated in the modulation of pain," the authors wrote.

Patients rated their mood, before and after the visceral distention, using the Stress Symptom Rating scale and they rated the intensity of their discomfort on a 3-point scale. In addition, they were all evaluated for depression and anxiety symptoms.

Overall, depression and anxiety scores fell within the normal range for all controls and 12 of the 14 IBS patients, but both scores were higher in IBS patients than controls, even when the two clinically elevated patients were excluded. Self-reported stress, anxiety, and anger were also higher in IBS patients. "The current results demonstrate that during certain expectation of experimental abdominal/pelvic discomfort, female IBS-C [IBS with constipation] patients are more anxious and less able than healthy controls to down-regulate activity within the CNS network activated by potentially aversive interoceptive stimuli," the authors noted. ■

ADVERSE REACTIONS

Clinical Trials Experience. The overall incidence of side effects reported in patients receiving sitagliptin and metformin was similar to that reported with patients receiving placebo and metformin.

In a 24-week placebo-controlled trial of sitagliptin 100 mg administered once daily added to a twice-daily metformin regimen, there were no adverse reactions reported regardless of investigator assessment of causality in ≥5% of patients and more commonly than in patients given placebo. Discontinuation of therapy due to clinical adverse reactions was similar to the placebo treatment group (sitagliptin and metformin, 1.9%; placebo and metformin, 2.5%).

The overall incidence of adverse reactions of hypoglycemia in patients treated with sitagliptin and metformin was similar to patients treated with placebo and metformin (100 mg sitagliptin and metformin, 1.3%; placebo and metformin, 2.1%). Adverse reactions of hypoglycemia were based on all reports of hypoglycemia; a concurrent glucose measurement was not required. The incidence of selected gastrointestinal adverse reactions in patients treated with sitagliptin and metformin was also similar to placebo and metformin: nausea (sitagliptin and metformin, 1.3%; placebo and metformin, 0.8%), vomiting (1.1%, 0.8%), abdominal pain (2.2%, 3.8%), and diarrhea (2.4%, 2.5%).

No clinically meaningful changes in vital signs or in ECG (including in QTc interval) were observed with the combination of sitagliptin and metformin.

The most common adverse experience in sitagliptin monotherapy reported regardless of investigator assessment of causality in ≥5% of patients and more commonly than in patients given placebo was nasopharyngitis.

The most common (>5%) established adverse reactions due to initiation of metformin therapy are diarrhea, nausea/vomiting, flatulence, abdominal discomfort, indigestion, asthenia, and headache.

Laboratory Tests

Sitagliptin. The incidence of laboratory adverse reactions was similar in patients treated with sitagliptin and metformin (7.6%) compared to patients treated with placebo and metformin (8.7%). In most but not all studies, a small increase in white blood cell count (approximately 200 cells/microL difference in WBC vs placebo; mean baseline WBC approximately 6600 cells/microL) was observed due to a small increase in neutrophils. This change in laboratory parameters is not considered to be clinically relevant.

Metformin hydrochloride. In controlled clinical trials of metformin of 29 weeks duration, a decrease to subnormal levels of previously normal serum Vitamin B₁₂ levels, without clinical manifestations, was observed in approximately 7% of patients. Such decrease, possibly due to interference with B₁₂ absorption from the B₁₂-intrinsic factor complex, is, however, very rarely associated with anemia and appears to be rapidly reversible with discontinuation of metformin or Vitamin B₁₂ supplementation [see Warnings and Precautions].

Postmarketing Experience. The following additional adverse reactions have been identified during postapproval use of JANUMET or sitagliptin, one of the components of JANUMET. Because these reactions are reported voluntarily from a population of uncertain size, it is generally not possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Hypersensitivity reactions include anaphylaxis, angioedema, rash, urticaria and exfoliative skin conditions including Stevens-Johnson syndrome [see Warnings and Precautions]; upper respiratory tract infection.

DRUG INTERACTIONS

Cationic Drugs. Cationic drugs (e.g., amiloride, digoxin, morphine, procainamide, quinidine, quinine, ranitidine, triamterene, trimethoprim, or vancomycin) that are eliminated by renal tubular secretion theoretically have the potential for interaction with metformin by competing for common renal tubular transport systems. Such interaction between metformin and oral cimetidine has been observed in normal healthy volunteers in both single- and multiple-dose metformin-cimetidine drug interaction studies, with a 60% increase in peak metformin plasma and whole blood concentrations and a 40% increase in plasma and whole blood metformin AUC. There was no change in elimination half-life in the single-dose study. Metformin had no effect on cimetidine pharmacokinetics. Although such interactions remain theoretical (except for cimetidine), careful patient monitoring and dose adjustment of JANUMET and/or the interfering drug is recommended in patients who are taking cationic medications that are excreted via the proximal renal tubular secretory system.

Digoxin. There was a slight increase in the area under the curve (AUC, 11%) and mean peak drug concentration (C_{max}, 18%) of digoxin with the coadministration of 100 mg sitagliptin for 10 days. These increases are not considered likely to be clinically meaningful. Digoxin, as a cationic drug, has the potential to compete with metformin for common renal tubular transport systems, thus affecting the serum concentrations of either digoxin, metformin or both. Patients receiving digoxin should be monitored appropriately. No dosage adjustment of digoxin or JANUMET is recommended.

Glyburide. In a single-dose interaction study in type 2 diabetes patients, coadministration of metformin and glyburide did not result in any changes in either metformin pharmacokinetics or pharmacodynamics. Decreases in glyburide AUC and C_{max} were observed, but were highly variable. The single-dose nature of this study and the lack of correlation between glyburide blood levels and pharmacodynamic effects make the clinical significance of this interaction uncertain.

Furosemide. A single-dose, metformin-furosemide drug interaction study in healthy subjects demonstrated that pharmacokinetic parameters of both compounds were affected by coadministration. Furosemide increased the metformin plasma and blood C_{max} by 22% and blood AUC by 15%, without any significant change in metformin renal clearance. When administered with metformin, the C_{max} and AUC of furosemide were 31% and 12% smaller, respectively, than when administered alone, and the terminal half-life was decreased by 32%, without any significant change in furosemide renal clearance. No information is available about the interaction of metformin and furosemide when coadministered chronically.

Nifedipine. A single-dose, metformin-nifedipine drug interaction study in normal healthy volunteers demonstrated that coadministration of nifedipine increased plasma metformin C_{max} and AUC by 20% and 9%, respectively, and increased the amount excreted in the urine. T_{max} and half-life were unaffected. Nifedipine appears to enhance the absorption of metformin. Metformin had minimal effects on nifedipine.

The Use of Metformin with Other Drugs. Certain drugs tend to produce hyperglycemia and may lead to loss of glycemic control. These drugs include the thiazides and other diuretics, corticosteroids, phenothiazines, thyroid products, estrogens, oral contraceptives, phenytoin, nicotinic acid, sympathomimetics, calcium channel blocking drugs, and isoniazid. When such drugs are administered to a patient receiving JANUMET the patient should be closely observed to maintain adequate glycemic control.

In healthy volunteers, the pharmacokinetics of metformin and propranolol, and metformin and ibuprofen were not affected when coadministered in single-dose interaction studies.

Metformin is negligibly bound to plasma proteins and is, therefore, less likely to interact with highly protein-bound drugs such as salicylates, sulfonamides, chloramphenicol, and probenecid, as compared to the sulfonylureas, which are extensively bound to serum proteins.

USE IN SPECIFIC POPULATIONS

Pregnancy

Pregnancy Category B.

JANUMET. There are no adequate and well-controlled studies in pregnant women with JANUMET or its individual components; therefore, the safety of JANUMET in pregnant women is not known. JANUMET should be used during pregnancy only if clearly needed.

Merck & Co., Inc. maintains a registry to monitor the pregnancy outcomes of women exposed to JANUMET while pregnant. Health care providers are encouraged to report any prenatal exposure to JANUMET by calling the Pregnancy Registry at (800) 986-8999.

No animal studies have been conducted with the combined products in JANUMET to evaluate effects on reproduction. The following data are based on findings in studies performed with sitagliptin or metformin individually.

Sitagliptin. Reproduction studies have been performed in rats and rabbits. Doses of sitagliptin up to 125 mg/kg (approximately 12 times the human exposure at the maximum recommended human dose) did not impair fertility or harm the fetus. There are, however, no adequate and well-controlled studies with sitagliptin in pregnant women.

Sitagliptin administered to pregnant female rats and rabbits from gestation day 6 to 20 (organogenesis) was not teratogenic at oral doses up to 250 mg/kg (rats) and 125 mg/kg (rabbits), or approximately 30 and 20 times human exposure at the maximum recommended human dose (MRHD) of 100 mg/day based on AUC comparisons. Higher doses increased the incidence of rib malformations in offspring at 1000 mg/kg, or approximately 100 times human exposure at the MRHD.

Sitagliptin administered to female rats from gestation day 6 to lactation day 21 decreased body weight in male and female offspring at 1000 mg/kg. No functional or behavioral toxicity was observed in offspring of rats.

Placental transfer of sitagliptin administered to pregnant rats was approximately 45% at 2 hours and 80% at 24 hours postdose. Placental transfer of sitagliptin administered to pregnant rabbits was approximately 66% at 2 hours and 30% at 24 hours.

Metformin hydrochloride. Metformin was not teratogenic in rats and rabbits at doses up to 600 mg/kg/day. This represents an exposure of about 2 and 6 times the maximum recommended human daily dose of 2000 mg based on body surface area comparisons for rats and rabbits, respectively. Determination of fetal concentrations demonstrated a partial placental barrier to metformin.

Nursing Mothers. No studies in lactating animals have been conducted with the combined components of JANUMET. In studies performed with the individual components, both sitagliptin and metformin are secreted in the milk of lactating rats. It is not known whether sitagliptin is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when JANUMET is administered to a nursing woman.

Pediatric Use. Safety and effectiveness of JANUMET in pediatric patients under 18 years have not been established.

Geriatric Use. **JANUMET.** Because sitagliptin and metformin are substantially excreted by the kidney and because aging can be associated with reduced renal function, JANUMET should be used with caution as age increases. Care should be taken in dose selection and should be based on careful and regular monitoring of renal function [see Warnings and Precautions].

Sitagliptin. Of the total number of subjects (N=3884) in Phase II and III clinical studies of sitagliptin, 725 patients were 65 years and over, while 61 patients were 75 years and over. No overall differences in safety or effectiveness were observed between subjects 65 years and over and younger subjects. While this and other reported clinical experience have not identified differences in responses between the elderly and younger patients, greater sensitivity of some older individuals cannot be ruled out.

Metformin hydrochloride. Controlled clinical studies of metformin did not include sufficient numbers of elderly patients to determine whether they respond differently from younger patients, although other reported clinical experience has not identified differences in responses between the elderly and young patients. Metformin should only be used in patients with normal renal function. The initial and maintenance dosing of metformin should be conservative in patients with advanced age, due to the potential for decreased renal function in this population. Any dose adjustment should be based on a careful assessment of renal function [see Contraindications, Warnings and Precautions].

Janumet
(sitagliptin/metformin HCl)
tablets



JANUMET is a trademark of Merck & Co., Inc. Copyright ©2008 Merck & Co., Inc. Whitehouse Station, NJ 08889, USA All rights reserved. 20803037(1)(105)-JMT