

Intellectual Impairment: Use Developmental Lens

BY DIANA MAHONEY
New England Bureau

BOSTON — Failure to use a developmental framework when assessing mental illness in children and adolescents with intellectual disabilities can hinder the accurate diagnosis of psychiatric disorders, and lead to inappropriate and possibly dangerous treatment, according to Lauren R. Charlot, Ph.D.

"It's important to consider the ways in which an individual's developmental stage affects his or her behavior and thoughts," Dr. Charlot said at the annual meeting of the American Society for Adolescent Psychiatry.

"Some symptoms or behaviors that would seem pathologic for a typically developing individual may be 'typical' for a person with a particular set of skills and challenges, especially when under stress," said Dr. Charlot, of the department of psychiatry at the University of Massachusetts, Worcester.

For example, she said, an adolescent with similar cognitive features to a typically developing preschool-age child might engage in self-talk—talking out loud to himself or herself—under stress. "Developmentally, this behavior is appropriate, but if you don't apply a developmental perspective, it seems abnormal and could easily be mistaken for psychotic ideation."

The most important consideration when evaluating individuals with intellectual delays is the link between developmental stage and cognitive processes. Studies have identified multiple develop-

mental effects on psychopathology in people with intellectual disabilities, Dr. Charlot noted. Among the developmental effects most likely to have an impact on psychiatric evaluation in this patient population are:

► **Magical thinking**, characterized by poor distinctions between fantasy and reality. "These kids may be talking to people who are not present, and they may be expressing wishes as beliefs," Dr. Charlot said. Unfortunately, she added, "this behavior is often misconstrued as hallucinations or delusions," even though it is a consequence of cognitive development rather than psychotic thinking.

► **Prelogical thinking**, which is characterized by primitive, incomplete thought processes and may present as seemingly tangential rambling. Normal among preschoolers, this mode of thought can be symptomatic of mental illness—particularly schizophrenia—in typically developing adolescents and adults, Dr. Charlot said. "This type of prelogical thinking is common at baseline in people with intellectual delays, and it tends to get exaggerated when they're under stress, which is often reported as 'racing thoughts.' But it's not a consequence of psychosis. It relates to an inability to understand logical relationships between subjects," she said.

► **Concrete thinking**, whereby language

and perceptions are interpreted literally. The limited ability to understand abstract language and ideas can lead individuals to misinterpret questions that are asked of them, which could be perceived as an indication of disturbed thought, Dr. Charlot said.

The clinical population of children and teens with developmental disabilities is heterogenous, so know what the individual's baseline behavior is.

► **Egocentrism**, or the tendency to assume that everyone shares one's own thoughts and the inability to understand the consequences of one's own behavior on others. In normally developing children, egocentric be-

havior begins to wane after toddler and preschool stages. Therefore, when it presents in adolescents, it can be misconstrued as narcissistic or oppositional behavior, according to Dr. Charlot.

To avoid the "trap" of perceiving developmentally driven behaviors as symptomatic of mental illness in individuals with intellectual disabilities, "You need to ask the question: 'what is usual behavior for this patient?'" Dr. Charlot stressed. "Because the clinical population is highly heterogenous, you need to know what the individual person's baseline [behavior] is. You cannot use general population reference group [data] to determine that a symptom or symptoms are evidence of psychosis or other psychopathology."

Additionally, Dr. Charlot continued, "ask yourself if the behaviors and symp-

toms are pathologic for a person functioning at this developmental level." For example, if you are assessing an adolescent who appears labile and "all over the place," she said, "that's not that unusual for an intellectually disabled individual who is experiencing stress."

Before assuming that alterations in mood and behavior are indicative of a psychiatric disorder in adolescents with intellectual impairment, "hunt for possible sources of physical distress," Dr. Charlot said.

Among the many medical problems that can manifest as behavioral issues are constipation, gastroesophageal reflux disease, seizure disorder, hypothyroidism, hypertension, anemia, candidiasis, urinary tract infection, diabetes, hypercholesterolemia, and obesity, she said, adding, "I can't tell you how many cases of adolescents with reported severe, aggressive behavior turned out to be constipated."

Often, because of poor language skills, these individuals are unable to articulate physical complaints, resulting in agitation, which may then be attributed to a psychiatric disorder, she said at the meeting cosponsored by the University of Texas at Dallas.

Finally, if and when a psychiatric syndrome is identified, "be careful not to automatically attribute any future alterations in mental status or behavior as being secondary to the psychiatric problem," Dr. Charlot said.

Under all circumstances, "comprehensive treatment should be directly derived from comprehensive assessment." ■

Early Family-Based Intervention Might Help Prevent Antisocial Behavior

BY CAROLYN SACHS
Contributing Writer

KAUI, HAWAII — Preschoolers at genetic risk for antisocial behavior may benefit from family-based preventive intervention, Dr. Glen O. Gabbard said at the annual meeting of the American College of Psychiatrists.

"I worked in the prison system for 6 years. One of the things you see again and again is that antisocial patients are not very responsive to individual therapy," he said.

If they don't receive any treatment until they are adults, "you're not going to get anywhere with them," said Dr. Gabbard, Brown Foundation Professor of Psychoanalysis and professor of psychiatry at Baylor College of Medicine, Houston. "We need to start thinking about early preventive approaches based on family therapy models."

In a recent study that he discussed, investigators enrolled 92 preschoolers who were considered

to be at substantial genetic risk for antisocial behavior because they had siblings with a history of juvenile delinquency (Arch. Gen. Psychiatry 2007;64:1172-9).

The children were randomized to one of two groups. In the fam-



Preventive family therapy might play a role in boosting activation of the amygdala.

DR. GABBARD

ily-intervention group, the preschoolers and their parents had 22 weekly group sessions and 10 biweekly home visits over a 6- to 8-month period. The control group received assessments and monthly telephone calls. Salivary cortisol levels were measured as an indicator of response to stress.

Since salivary cortisol levels are lower in people who exhibit antisocial behavior and have conduct

problems, "the thought here is that maybe the early-life experience can alter cortisol release," he noted.

Compared with the control group, children who had undergone family-based intervention had increased salivary cortisol levels when faced with entry into an unfamiliar group of peers. "Something changed in their reading of social groups so that they had more anxiety," Dr. Gabbard said.

"The antisocial patient doesn't engage the amygdala in the way that a borderline patient does," he observed. The antisocial patient has "less of a fear response." Family-based preventive intervention may play a role in boosting activation of the amygdala and modifying the hypothalamic-pituitary-adrenal axis when someone is faced with a threatening situation, he said.

"Preventive family therapy may give us some hope in what has been a rather dismal history of treating the psychopathic or antisocial patient," said Dr. Gabbard, who disclosed no relevant conflicts. ■

Factors Identified for Earlier Dx of ASD

BY KERRI WACHTER
Senior Writer

PHILADELPHIA — Children are likely to be diagnosed with autism spectrum disorders at a younger age if there is a shorter period between referral and evaluation, Dr. Ginger Janow said.

Early diagnosis of autism spectrum disorders can make a big difference in outcomes.

"The one thing that's been shown consistently [to] affect outcomes is early intensive behavioral intervention," said Dr. Janow of the Children's Hospital at Montefiore, New York, at the annual meeting of the Eastern Pediatric Research Society. The effectiveness of these interventions depends on the age at which they are started.

The researchers performed a retrospective chart review of 116 children diagnosed with autism spectrum disorder

(ASD) at the Seaver and New York Autism Center of Excellence at the Mount Sinai School of Medicine, New York, between 1995 and 2005. Specialists at the center conduct initial evaluations for ASD.

Supplementary evaluations are performed when additional assessment is warranted. The average age of diagnosis at the center was 27 months, and the average time between referral and evaluation was 2 months.

An analysis showed that earlier age of diagnosis was correlated with diagnosis at the primary evaluation (rather than at a supplemental evaluation), decreased time between referral and evaluation, and increased fine-motor and adaptive delays.

Early diagnosis was not correlated with insurance status, estimated income, or language delays. ■