

Response to Abatacept May Increase Over Time

BY NANCY WALSH
New York Bureau

PARIS — Patients with rheumatoid arthritis receiving abatacept showed an increasing magnitude of clinical response between months 6 and 12, Dr. Michael Schiff reported.

In RA, it's important to evaluate efficacy not only at the group level—mean changes that occur in a group as a whole over time—but also at the individual level, and to measure changes in a patient's response over time, Dr. Schiff said in a press conference at the annual European Congress of Rheumatology.

In a post-hoc analysis of ATTEST (Abatacept or infliximab versus placebo, a Trial for Tolerability, Efficacy and Safety in Treating RA), patients who achieved an ACR 20 at 6 months were reevaluated at 12 months to determine whether they further improved, retained, or lost their response.

The larger ATTEST trial, sponsored by Bristol-Myers Squibb Co., included 431 patients who had an inadequate response to methotrexate, and randomized them to abatacept, approximately 10 mg/kg every 4 weeks; infliximab, 3 mg/kg every 8 weeks; or placebo. They also were on background methotrexate, 10-30 mg/week.

At baseline, patients' mean age was 49 years and mean disease duration was 8 years. The majority were women. Mean number of swollen joints was 20, mean number of tender joints was 30, and mean disease activity score (DAS) 28 was 6.4.

At month 6, a total of 32 patients receiving abatacept and 27 patients receiving infliximab had achieved an ACR 20 response, though not ACR 50 or ARC 70 responses. The responses in this subgroup of patients were then analyzed at 12 months.

A similar percentage of patients receiving

abatacept and infliximab achieved at least an ACR 50 response at 1 year. A total of 28.1% of patients on abatacept and 29.6% of patients on infliximab continued to improve to this degree, a clinically significant response, said Dr. Schiff, a clinical professor of medicine in the rheumatology division, University of Colorado, Denver.

The proportion of patients who maintained an ACR 20 response at 1 year was 59.4% for abatacept and 44.4% for infliximab, while the proportions who lost their

ACR 20 response at 1 year were 12.5% and 25.9%, respectively.

Among the 24 patients treated with abatacept who achieved a low disease activity score (DAS28 below 3.2) at 6 months, 41.7% went on to clinical remission (DAS28 below 2.6) at 1 year. Among the 25 patients treated with infliximab who achieved a low disease activity score at 6 months, 28% went on to remission at 1 year.

A total of 12.5% and 16% of the abatacept- and infliximab-treated patients re-

tained their low disease activity score at 1 year, while 45.8% and 56% lost this response by 1 year.

"These findings highlight the importance of examining clinical benefits at the level of the individual patient," Dr. Schiff wrote in a poster. "Abatacept has the potential to provide clinical benefits to patients that increase in magnitude over time."

Dr. Schiff has received research grants and consulting fees from Bristol-Myers Squibb and Centocor. ■

Sacroiliitis, PsA Often Linked

PARIS — Sacroiliitis is an often-underappreciated hallmark of psoriatic arthritis, according to Dr. Augustin Sellas of Vall d'Hebron Hospital, Barcelona.

Peripheral joint disease in psoriatic arthritis patients draws considerable physician attention.

But sacroiliitis is highly prevalent as well, he noted at the annual European Congress of Rheumatology.

Dr. Sellas retrospectively reviewed x-rays of 128 PsA patients, 51% of whom had radiographic evidence of sacroiliitis.

The prevalence was similar in men and women. Sacroiliitis was bilateral in 47 of 65 affected patients, or 72%.

Seventeen percent of psoriatic arthritis patients were HLA-B27 positive. Sacroiliitis was present in three-quarters of this subgroup, with all but one case being bilateral.

The reported prevalence of sacroiliitis in PsA patients has ranged from 30% to 78% in other studies, said Dr. Sellas.

—Bruce Jancin

Bisphosphonate therapy can't hold up without calcium & vitamin D^{1†}

Yet 90% of women may have low calcium intake³

ONLY OS-CAL is proven to help reduce the risk of hip fractures by 29%*²

OS-CAL. Helping to support your patient's bisphosphonate therapy

- > Bisphosphonates require adequate calcium to be effective^{1†}
- > Demonstrated **29%** hip fracture risk reduction*(NIH Study, n=21,406)²
- > Improved bone mineral density and strength (n=1,731)⁴
- > Proven effective in more clinical studies than any other calcium supplement
- > Improved BMD has been shown to be independent of bisphosphonate therapy^{2†}
- > Unsurpassed calcium absorption^{5,6}



OS-CAL
calcium carbonate + vitamin D₃

An essential bisphosphonate adjunct
Don't forget the OS-CAL twice a day every day with meals

* Based on NIH study, when taken as directed, OS-CAL formulation of 500 mg calcium + 200 IU vitamin D₃.

† Prescribing information for: Fosamax® & Fosamax plus D™ - Merck, Actonel® & Actonel® with Calcium - Procter & Gamble, Boniva® - Roche
1. Guenther C et al. Nothing works without Calcium and Vitamin D. Oasis online abstracts Presentation T393. 2. Jackson RD et al. Calcium plus Vitamin D supplementation and the Risk of Fractures. *N Engl J Med* V354;7 Feb 16, 2006. 3. USDA's Continuing Survey of Food Intakes by Individuals 1994-1996. 4. Chen Z et al. The effect of Calcium plus Vitamin D Supplement on Hip Geometric Structures: Results from the Women's Health Initiative CaD Trial. Oasis online abstracts Presentation T1207. 5. Recker RR. Calcium absorption and achlorhydria. *N Engl J Med*. 1985;313:70-73. 6. Bo-Linn GW, et al. An evaluation of the importance of gastric acid secretion in the absorption of dietary calcium. *J Clin Invest*. 1984;73:640-657. © GlaxoSmithKline 2008