

Ongoing Trial Compares Anorexia Therapies

BY SUSAN LONDON
Contributing Writer

SEATTLE — The relative efficacy of three treatments for anorexia nervosa appears to shift with long-term follow-up according to the results of an ongoing analysis of data from a randomized, controlled trial.

The treatment that was the most efficacious at the end of therapy appeared to be the least so at 5 years. But differences among the therapies were much less sig-

nificant at that point, Virginia V.W. McIntosh, Ph.D., reported at an international conference sponsored by the Academy for Eating Disorders.

Dr. McIntosh, a senior clinical psychologist at the University of Otago, Christchurch, New Zealand, described ongoing analyses of data from a randomized, controlled trial that compared three treatments—cognitive-behavioral therapy (CBT), interpersonal psychotherapy (IPT), and specialist supportive clinical manage-

ment (SSCM)—among 56 patients with anorexia nervosa.

End-of-treatment results showed that among the 35 patients who completed all sessions, SSCM was superior to both CBT and IPT in terms of global anorexia nervosa status (*Am. J. Psychiatry* 2005; 162:741-7).

One of the new analyses focused on therapist adherence to the protocol for a specific treatment.

Dr. McIntosh and her colleagues measured adherence with a modified version of the Collaborative Study Psychotherapy Rating Scale, which had 28 items unique to IPT, 27



unique to CBT, and 3 unique to SSCM. "I think [adherence] speaks to the distinctiveness of CBT and SSCM, which is important here," Dr. McIntosh said at the meeting, which was cosponsored by the University of New Mexico.

Another new, ongoing analysis of the trial data focused on long-term outcomes. Data at 5 years were available for 45 patients (80% of those initially randomized), Dr. McIntosh said. "The differences at 5 years are much less than the differences at end of treatment," she reported.

"What we see is that the SSCM group has lost some ground in terms of the proportion with a good outcome, CBT has gained some ground at about the same rate at which SSCM has lost ground, and IPT has gained even more ground," she said.

As a result, the proportion of patients with a good outcome at the 5-year time point was highest for IPT, intermediate for CBT, and lowest for SSCM.

For example, the CBT subscale scores were highest for CBT sessions. In contrast, ratings for the therapy-nonspecific subscale did not differ depending on which therapy the patient had received.

For example, the CBT subscale scores were highest for CBT sessions. In contrast, ratings for the therapy-nonspecific subscale did not differ depending on which therapy the patient had received.

For example, the CBT subscale scores were highest for CBT sessions. In contrast, ratings for the therapy-nonspecific subscale did not differ depending on which therapy the patient had received.

For example, the CBT subscale scores were highest for CBT sessions. In contrast, ratings for the therapy-nonspecific subscale did not differ depending on which therapy the patient had received.

LEXAPRO® (escitalopram oxalate) TABLETS/ORAL SOLUTION

(5% and 4%); Fatigue (5% and 2%). **Psychiatric Disorders:** Insomnia (9% and 4%); Somnolence (6% and 2%); Appetite Decreased (3% and 1%); Libido Decreased (3% and 1%). **Respiratory System Disorders:** Rhinitis (5% and 4%); Sinusitis (3% and 2%). **Urogenital:** Ejaculation Disorder* (9% and <1%); Impotence* (3% and <1%); Anorgasmia* (2% and <1%). *Events reported by at least 2% of patients treated with Lexapro are reported, except for the following events which had an incidence on placebo - Lexapro: headache, upper respiratory tract infection, back pain, pharyngitis, inflicted injury, anxiety. †Primarily ejaculatory delay. ‡Denominator used was for males only (N=225 Lexapro; N=188 placebo). †Denominator used was for females only (N=490 Lexapro; N=404 placebo). **Generalized Anxiety Disorder Table 3** numerates the incidence, rounded to the nearest percent of treatment-emergent adverse events that occurred among 429 GAD patients who received Lexapro 10 to 20 mg/day in placebo-controlled trials. Events included are those occurring in 2% or more of patients treated with Lexapro and for which the incidence in patients treated with Lexapro was greater than the incidence in placebo-treated patients. The most commonly observed adverse events in Lexapro patients (incidence of approximately 5% or greater and approximately twice the incidence in placebo-treated patients) were nausea, ejaculation disorder (primarily ejaculatory delay), insomnia, fatigue, decreased libido, and anorgasmia (see TABLE 3). **TABLE 3: Treatment-Emergent Adverse Events: Incidence in Placebo-Controlled Clinical Trials for Generalized Anxiety Disorder* (Percentage of Patients Reporting Event) Body System/Adverse Event (Lexapro (N=429) and Placebo (N=427)).** **Autonomic Nervous System Disorders:** Dry Mouth (9% and 5%); Sweating Increased (4% and 1%). **Central & Peripheral Nervous System Disorders:** Headache (24% and 17%); Paresthesia (2% and 1%). **Gastrointestinal Disorders:** Nausea (18% and 8%); Diarrhea (8% and 6%); Constipation (5% and 4%); Indigestion (3% and 2%); Vomiting (3% and 1%); Abdominal Pain (2% and 1%); Flatulence (2% and 1%); Toothache (2% and 0%). **General:** Fatigue (8% and 2%); Influenza-like symptoms (5% and 4%). **Musculoskeletal:** Neck/Shoulder Pain (3% and 1%). **Psychiatric Disorders:** Somnolence (13% and 7%); Insomnia (12% and 6%); Libido Decreased (7% and 2%); Dreaming Abnormal (3% and 2%); Appetite Decreased (3% and 1%); Lethargy (3% and 1%); Yawning (2% and 1%). **Urogenital:** Ejaculation Disorder* (14% and 2%); Anorgasmia† (6% and <1%); Menstrual Disorder (2% and 1%). *Events reported by at least 2% of patients treated with Lexapro are reported, except for the following events which had an incidence on placebo - Lexapro: inflicted injury, dizziness, back pain, upper respiratory tract infection, rhinitis, pharyngitis. †Primarily ejaculatory delay. ‡Denominator used was for males only (N=182 Lexapro; N=195 placebo). †Denominator used was for females only (N=247 Lexapro; N=232 placebo). **Dose Dependency of Adverse Events** The potential dose dependency of common adverse events (defined as an incidence rate of ≥ 5% in either the 10 mg or 20 mg Lexapro groups) was examined on the basis of the combined incidence of adverse events in two fixed-dose trials. The overall incidence rates of adverse events in 10 mg Lexapro-treated patients (66%) was similar to that of the placebo-treated patients (61%), while the incidence rate in 20 mg/day Lexapro-treated patients was greater (86%). **Table 4** shows common adverse events that occurred in the 20 mg/day Lexapro group with an incidence that was approximately twice that of the 10 mg/day Lexapro group and approximately twice that of the placebo group. **TABLE 4: Incidence of Common Adverse Events* in Patients with Major Depressive Disorder Receiving Placebo (N=311), 10 mg/day Lexapro (N=310), 20 mg/day Lexapro (N=125).** **Insomnia** (4%, 7%, 14%); **Diarrhea** (5%, 6%, 14%); **Dry Mouth** (3%, 4%, 9%); **Somnolence** (1%, 4%, 9%); **Dizziness** (2%, 4%, 7%); **Sweating Increased** (<1%, 3%, 8%); **Constipation** (1%, 3%, 6%); **Fatigue** (2%, 2%, 6%); **Indigestion** (1%, 2%, 6%). *Adverse events with an incidence rate of at least 5% in either the Lexapro groups and with an incidence rate in the 20 mg/day Lexapro group that was approximately twice that of the 10 mg/day Lexapro group and the placebo group. **Male and Female Sexual Dysfunction with SSRIs** Although changes in sexual desire, sexual performance, and sexual satisfaction often occur as manifestations of a psychiatric disorder, they may also be a consequence of pharmacologic treatment. In particular, some evidence suggests that SSRIs can cause such untoward sexual experiences. Reliable estimates of the incidence and severity of untoward experiences involving sexual desire, performance, and satisfaction are difficult to obtain, however, in part because patients and physicians may be reluctant to discuss them. Accordingly, estimates of the incidence of untoward sexual experience and performance cited in product labeling are likely to underestimate their actual incidence. **Table 5** shows the incidence rates of sexual side effects in patients with major depressive disorder and GAD in placebo-controlled trials. **TABLE 5: Incidence of Sexual Side Effects in Placebo-Controlled Clinical Trials (In Males Only: Adverse Event: Lexapro (N=407) and Placebo (N=383)).** Ejaculation Disorder (primarily ejaculatory delay) (12% and 1%); Libido Decreased (6% and 2%); Impotence (2% and <1%). (In Females Only: Lexapro (N=373) and Placebo (N=356)). Libido Decreased (3% and 1%); Anorgasmia (3% and <1%). There are no adequately designed studies examining sexual dysfunction with escitalopram treatment. Priapism has been reported with all SSRIs. While it is difficult to know the precise risk of sexual dysfunction associated with the use of SSRIs, physicians should routinely inquire about such possible side effects. **Vital Sign Changes** Lexapro and placebo groups were compared with respect to (1) mean change from baseline in vital signs (pulse, systolic blood pressure, and diastolic blood pressure) and (2) the incidence of patients meeting criteria for potentially clinically significant changes from baseline in these variables. These analyses did not reveal any clinically important changes in vital signs associated with Lexapro treatment. In addition, a comparison of supine and standing vital sign measures in subjects receiving Lexapro indicated that Lexapro treatment is not associated with orthostatic changes. **Weight Changes** Patients treated with Lexapro in controlled trials did not differ from placebo-treated patients with regard to clinically important change in body weight. **Laboratory Changes** Lexapro and placebo groups were compared with respect to (1) mean change from baseline in various serum chemistry, hematology, and urinalysis variables, and (2) the incidence of patients meeting criteria for potentially clinically significant changes from baseline in these variables. These analyses revealed no clinically important changes in laboratory test parameters associated with Lexapro treatment. **ECG Changes** Electrocardiograms from Lexapro (N=625), racemic citalopram (N=351), and placebo (N=527) groups were compared with respect to (1) mean change from baseline in various ECG parameters and (2) the incidence of patients meeting criteria for potentially clinically significant changes from baseline in these variables. These analyses revealed (1) a decrease in heart rate of 2.2 bpm for Lexapro and 2.7 bpm for racemic citalopram, compared to an increase of 0.3 bpm for placebo and (2) an increase in QTc interval of 3.9 msec for Lexapro and 3.7 msec for racemic citalopram, compared to 0.5 msec for placebo. Neither Lexapro nor racemic citalopram were associated with the development of clinically significant ECG abnormalities. **Other Events Observed During the Premarketing Evaluation of Lexapro** Following is a list of WHO terms that reflect treatment-emergent adverse events, as defined in the introduction to the ADVERSE REACTIONS section, reported by the 1428 patients treated with Lexapro for periods of up to one year in double-blind or open-label clinical trials during its premarketing evaluation. All reported events are included except those already listed in Tables 2 & 3, those occurring in only one patient, event terms that are so general as to be uninformative, and those that are unlikely to be drug related. It is important to emphasize that, although the events reported occurred during treatment with Lexapro, they were not necessarily caused by it. Events are further categorized by body system and listed in order of decreasing frequency according to the following definitions: frequent adverse events are those occurring on one or more occasions in at least 1/100 patients; infrequent adverse events are those occurring in less than 1/100 patients but at least 1/1000 patients. **Cardiovascular - Frequent:** palpitation, hypertension. **Infrequent:** bradycardia, tachycardia, ECG abnormal, flushing, varicose vein. **Central and Peripheral Nervous System Disorders - Frequent:** light-headed feeling, migraine. **Infrequent:** tremor, vertigo, restless legs, shaking, twitching, dysequilibrium, tics, carpal tunnel syndrome, muscle contractions involuntary, sluggishness, coordination abnormal, faintness, hyperreflexia, muscular tone increased. **Gastrointestinal Disorders - Frequent:** heartburn, abdominal cramp, gastroenteritis. **Infrequent:** gastroesophageal reflux, bloating, abdominal discomfort, dyspepsia, increased stool frequency, belching, gastritis, hemorroids, gagging, polyposis gastric, swallowing difficult. **General - Frequent:** allergy, pain in limb, fever, hot flashes, chest pain. **Infrequent:** edema of extremities, chills, tightness of chest, leg pain, asthma, syncope, malaise, anaphylaxis, fall. **Hemic and Lymphatic Disorders - Frequent:** bruise, anemia, nosebleed, hematoma, lymphadenopathy cervical. **Metabolic and Nutritional Disorders - Frequent:** increased weight. **Infrequent:** decreased weight, hyperglycemia, thirst, bilirubin increased, hepatic enzymes increased, gout, hypercholesterolemia. **Musculoskeletal System Disorders - Frequent:** arthralgia, myalgia. **Infrequent:** jaw stiffness, muscle cramp, muscle stiffness, arthritis, muscle weakness, back discomfort, arthropathy, jaw pain, joint stiffness. **Psychiatric Disorders - Frequent:** appetite increased, lethargy, irritability, concentration impaired. **Infrequent:** jitteriness, panic reaction, agitation, apathy, forgetfulness, depression aggravated, nervousness, restlessness aggravated, suicide attempt, amnesia, anxiety attack, bruising, carbohydrate craving, confusion, depersonalization, disorientation, emotional lability, feeling unreal, tremulousness nervous, crying abnormal, depression, excitability, auditory hallucination, suicidal tendency. **Reproductive Disorders/Female* - Frequent:** menstrual cramps, menstrual disorder. **Infrequent:** menorrhagia, breast neoplasm, pelvic inflammation, premenstrual syndrome, spotting between menses. *% based on female subjects only. **N=905 Respiratory System Disorders - Frequent:** bronchitis, sinus congestion, coughing, nasal congestion, sinus headache. **Infrequent:** asthma, breath shortness, laryngitis, pneumonia, tracheitis. **Skin and Appendages Disorders - Frequent:** rash. **Infrequent:** pruritus, acne, alopecia, eczema, dermatitis, dry skin, folliculitis, lipoma, furunculosis, dry lips, skin nodule. **Special Senses - Frequent:** vision blurred, tinnitus. **Infrequent:** taste alteration, earache, conjunctivitis, vision abnormal, dry eyes, eye irritation, visual disturbance, eye infection, pupils dilated, metallic taste. **Urinary System Disorders - Frequent:** urinary frequency, urinary tract infection. **Infrequent:** urinary urgency, kidney stone, dysuria, blood in urine. **Events Reported Subsequent to the Marketing of Escitalopram** - Although no causal relationship to escitalopram treatment has been found, the following adverse events have been reported to have occurred in patients and to be temporally associated with escitalopram treatment during post marketing spontaneous and clinical trial experience and were not observed during the premarketing evaluation of escitalopram: Blood and Lymphatic System Disorders: hemolytic anemia, leukopenia, thrombocytopenia. Cardiac Disorders: atrial fibrillation, cardiac failure, myocardial infarction, torsade de pointes, ventricular arrhythmia, ventricular tachycardia. Endocrine Disorders: diabetes mellitus, hyperprolactinemia, SIADH. Eye Disorders: diplopia, glaucoma. Gastrointestinal Disorders: gastrointestinal hemorrhage, pancreatitis, rectal hemorrhage. General Disorders and Administration Site Conditions: abnormal gait. Hepatobiliary Disorders: fulminant hepatitis, hepatic failure, hepatic necrosis, hepatitis. Immune System Disorders: allergic reaction. Investigations: electrocardiogram QT prolongation, INR increased, prothrombin decreased. Metabolism and Nutrition Disorders: hypoglycemia, hypokalemia. Musculoskeletal and Connective Tissue Disorders: rhabdomyolysis. Nervous System Disorders: akathisia, choreoathetosis, dysarthria, dyskinesia, dystonia, extrapyramidal disorders, grand mal seizures (or convulsions), hypoaesthesia, myoclonus, neuroleptic malignant syndrome, nystagmus, seizures, serotonin syndrome, tardive dyskinesia. Pregnancy, Puerperium and Perinatal Conditions: spontaneous abortion. Psychiatric Disorders: acute psychosis, aggression, anger, delirium, delusion, nightmare, paranoia, visual hallucinations. Renal and Urinary Disorders: acute renal failure. Reproductive System and Breast Disorders: priapism. Respiratory, Thoracic and Mediastinal Disorders: pulmonary embolism. Skin and Subcutaneous Tissue Disorders: angioedema, ecchymosis, erythema multiforme, photosensitivity reaction, Stevens Johnson Syndrome, toxic epidermal necrolysis, urticaria. Vascular Disorders: deep vein thrombosis, hypotension, orthostatic hypotension, phlebitis thrombosis. Forest Pharmaceuticals, Inc. Subsidiary of Forest Laboratories, Inc. St. Louis, MO 63045 USA Licensed from H. Lundbeck A/S Rev. 04/08 02008 Forest Laboratories, Inc.

Use of Zolpidem Appears Safe During Pregnancy

BY KERRI WACHTER
Senior Writer

WASHINGTON — Even though the sleeping aid zolpidem does cross the placenta, use of the drug during pregnancy does not appear to significantly affect outcomes, a study of 45 women shows.

The study, presented as a poster at the annual meeting of the American Psychiatric Association, included pregnant women who were enrolled in a prospective study of the pharmacokinetics of psychotropic drugs during pregnancy and who were treated with zolpidem (Ambien) during pregnancy. Maternal diagnoses were determined using the Structured Clinical Interview for DSM-IV (SCID). Maternal and cord blood were obtained at delivery when possible.

The placental passage rate was calculated as the ratio of medication concentration in the umbilical cord plasma to that in maternal plasma. When umbilical cord concentrations were below the limit of detection (less than 4.0 ng/mL), this value was used for data analysis. This approach was thought to be conservative, erring toward overestimation of fetal exposure to zolpidem. When both maternal and umbilical plasma concentrations were less

than the detection limit, the pair was excluded from the analysis.

Obstetrical and neonatal outcomes among women who had given birth to a live infant after taking zolpidem during pregnancy were compared with outcomes among a group of 45 women who were matched for age, race, level of education, SCID diagnosis, and pregnancy exposure to the same classes of psychotropics.

For women who took zolpidem during pregnancy, exposure by trimester included 38% in the first trimester, 56% in the second trimester, and 38% in the third trimester. The average zolpidem exposure during pregnancy was 14 weeks, and the average dose was 9 mg.

No statistically significant differences were found between the two groups in terms of obstetrical and neonatal outcomes. However, a trend toward preterm delivery and low-birth-weight infants was seen among women on zolpidem during pregnancy. "It is unclear if these outcomes were driven by zolpidem exposure and/or sleep disturbance or other pharmacological intervention in pregnancy," wrote Sandra Juric and her colleagues at Emory University's Women's Mental Health Program in Atlanta. Ms. Juric reported no conflicts of interest.