

# Insurance Not a Barrier for Most Patients in ED

BY SHERRY BOSCHERT  
San Francisco Bureau

SAN FRANCISCO — Most adults seen in emergency departments have health insurance and a usual source of care besides the ED, countering the common notion that uninsured patients are clogging up emergency departments because they have nowhere else to go, Ellen J. Weber, M.D., said at the annual meeting of the American College of Emergency Physicians. (See box on p. 73.)

Using data on a population-based sample of 49,603 adults, she and her associates estimated that more than 45 million U.S. adults made nearly 80 million visits to EDs during a recent 1-year period. Of those who went to emergency departments, 83% said they had a usual source of care other than the ED (accounting for 82% of ED visits), and 85% had some form of health

**Patients who used the ED for their usual care or who had no source of usual care were responsible for only 17% of ED visits in the study.**

insurance (accounting for 85% of ED visits), she said. Improving the delivery of outpatient care may be the key to decreasing use of the ED, the investigators concluded. The study—one of a few to compare ED users with non-

users in these respects—showed that uninsured adults were no more likely to visit the ED during the study period than were people with private insurance. Compared with privately insured patients, those with Medicaid or Medicare were 51% and 19% more likely, respectively, to use the ED. Adults with no usual source of care were 25% less likely to visit the ED than were adults whose usual care came from a private physician. Patients who used the ED for their usual care or who had no source of usual care were responsible for only 17% of ED visits.

People who used the ED most had poor health or changes in their usual care or insurance, the study found.

## Resources Offer Medicare Help

The Medicare Rights Center offers three weekly newsletters that provide news and information on health care issues for older adults and people with disabilities.

All three of the newsletters—"Dear Marci," "Asclepius," and "Medicare Watch"—are available free of charge. For more information about the three publications, visit [www.medicarerights.org/periodicalsframeset.html](http://www.medicarerights.org/periodicalsframeset.html).

Also, AARP now offers Spanish-speaking seniors an online resource for information about Medicare's new drug benefit. The site is located at [www.aarpsegundajuventud.org/spanish/drugs](http://www.aarpsegundajuventud.org/spanish/drugs).

People with poor physical health were more than twice as likely to have visited an ED as were people in good health, and they accounted for nearly half of all ED visits. People with poor mental health were 51% more likely to go to an ED than were people with good mental health. People who had had a change in their usual source of care were 30% more likely to seek help in an ED than were people with stable care. A change in insurance increased the odds of an ED visit by 15%. Poverty increased

the likelihood of an ED visit, but patients with incomes below the poverty level accounted for only one in five visits.

"People affected by ED overcrowding and closures of emergency departments are the seriously ill," said Dr. Weber, medical director of the emergency department at the University of California, San Francisco.

Data for the study came from the Community Tracking Study Household Survey. The Center for Studying Health System

Change, a nonpartisan policy research group in Washington, conducted the survey. The survey data, from July 2000 through June 2001, included interviews in English or Spanish with up to eight adults per household in 60 communities and in a national sample, plus administration of the SF-12 Health Survey.

Several factors often thought to decrease the use of EDs were not associated with fewer ED visits, including enrollment in an HMO, early availability of

*When insomnia adds to your patients' problems*

## A good night's sleep...



### AMBIEN can provide a good night's sleep

#### AMBIEN treats all 3 symptoms of insomnia

Trouble falling asleep	AMBIEN decreases time to fall asleep vs placebo <sup>1</sup>
Trouble staying asleep	AMBIEN reduces number of awakenings vs placebo <sup>2</sup>
Waking too early	AMBIEN increases total sleep time vs placebo <sup>3</sup>

AMBIEN is indicated for the short-term treatment of insomnia. In elderly or debilitated patients, or patients with hepatic dysfunction, treatment should be initiated with a 5-mg dose and patients closely monitored. Due to its rapid onset of action, patients should take AMBIEN right before going to bed and when ready for sleep. Patients should not take AMBIEN unless they are prepared to get a full night's sleep (7 to 8 hours) to avoid residual effects. Until they know how it will affect their physical or mental performance upon awakening, patients should not drive or operate hazardous machinery after taking AMBIEN or any other sleep medication. During short-term treatment with AMBIEN, the most commonly observed adverse effects in controlled clinical trials were drowsiness (2%), dizziness (1%), and diarrhea (1%). Because individuals with a history of addiction or substance abuse are at increased risk of habituation and dependence, they should be under careful surveillance when receiving AMBIEN or any other hypnotic. AMBIEN is classified as a Schedule IV controlled substance. Sedative hypnotics have produced withdrawal signs and symptoms following abrupt discontinuation. Hypnotics should generally be limited to 7 to 10 days of use, and reevaluation of the patient is recommended if they are taken for more than 2 to 3 weeks. Prescriptions for AMBIEN should not exceed a 1-month supply.

For more information, visit our Web site at [www.ambien.com](http://www.ambien.com)

©2004 Sanofi-Synthelabo Inc.