

Adverse Event	amlodipine		Placebo	
	M=% (N=1218)	F=% (N=512)	M=% (N=914)	F=% (N=336)
Edema	5.6	14.6	1.4	5.1
Flushing	1.5	4.5	0.3	0.9
Palpitations	1.4	3.3	0.9	0.9
Somnolence	1.3	1.6	0.8	0.3

The following events occurred in $\leq 1\%$ but $>0.1\%$ of patients treated with amlodipine in controlled clinical trials or under conditions of open trials or marketing experience where a causal relationship is uncertain; they are listed to alert the physician to a possible relationship: **Cardiovascular:** arrhythmia (including ventricular tachycardia and atrial fibrillation), bradycardia, chest pain, hypotension, peripheral ischemia, syncope, tachycardia, postural dizziness, postural hypotension, vasculitis. **Central and Peripheral Nervous System:** hypoesthesia, neuropathy peripheral, paresthesia, tremor, vertigo. **Gastrointestinal:** anorexia, constipation, dyspepsia, dysphagia, diarrhea, flatulence, pancreatitis, vomiting, gingival hyperplasia. **General:** allergic reaction, asthenia, back pain, hot flushes, malaise, pain, rigors, weight gain, weight decrease. **Musculoskeletal System:** arthralgia, arthrosis, muscle cramps, myalgia. **Psychiatric:** sexual dysfunction (male and female), insomnia, nervousness, depression, abnormal dreams, anxiety, depersonalization. **Respiratory System:** dyspnea, epistaxis. **Skin and Appendages:** angioedema, erythema multiforme, pruritus, rash, rash erythematous, rash maculopapular. **Special Senses:** abnormal vision, conjunctivitis, diplopia, eye pain, tinnitus. **Urinary System:** micturition frequency, micturition disorder, nocturia. **Autonomic Nervous System:** dry mouth, sweating increased. **Metabolic and Nutritional:** hyperglycemia, thirst. **Hematologic:** leukopenia, purpura, thrombocytopenia. The following events occurred in $\leq 0.1\%$ of patients treated with amlodipine in controlled clinical trials or under conditions of open trials or marketing experience: cardiac failure, pulse irregularity, extrasystoles, skin discoloration, urticaria, skin dryness, alopecia, dermatitis, muscle weakness, twitching, ataxia, hypertonia, migraine, cold and clammy skin, apathy, agitation, amnesia, gastritis, increased appetite, loose stools, coughing, rhinitis, dysuria, polyuria, parosmia, taste perversion, abnormal visual accommodation, and xerophthalmia. Other reactions occurred sporadically and cannot be distinguished from medications or concurrent disease states such as myocardial infarction and angina. Amlodipine therapy has not been associated with clinically significant changes in routine laboratory tests. No clinically relevant changes were noted in serum potassium, serum glucose, total triglycerides, total cholesterol, HDL cholesterol, uric acid, blood urea nitrogen, or creatinine. The following postmarketing event has been reported infrequently with amlodipine treatment where a causal relationship is uncertain: gynecomastia. In postmarketing experience, jaundice and hepatic enzyme elevations (mostly consistent with cholestasis or hepatitis) in some cases severe enough to require hospitalization have been reported in association with use of amlodipine. Amlodipine has been used safely in patients with chronic obstructive pulmonary disease, well-compensated congestive heart failure, peripheral vascular disease, diabetes mellitus, and abnormal lipid profiles. **The Atorvastatin Component of CADUET:** Atorvastatin is generally well-tolerated. Adverse reactions have usually been mild and transient. In controlled clinical studies of 2502 patients, $<2\%$ of patients were discontinued due to adverse experiences attributable to atorvastatin calcium. The most frequent adverse events thought to be related to atorvastatin calcium were constipation, flatulence, dyspepsia, and abdominal pain. Clinical Adverse Experiences: Adverse experiences reported in $\geq 2\%$ of patients in placebo-controlled clinical studies of atorvastatin, regardless of causality assessment, are shown in Table 3.

Table 3. Adverse Events in Placebo-Controlled Studies (% of Patients)

Body System/ Adverse Event	Placebo N=270	atorvastatin			
		10 mg N=863	20 mg N=36	40 mg N=79	80 mg N=94
BODY AS A WHOLE					
Infection	10.0	10.3	2.8	10.1	7.4
Headache	7.0	5.4	16.7	2.5	6.4
Accidental Injury	3.7	4.2	0.0	1.3	3.2
Flu Syndrome	1.9	2.2	0.0	2.5	3.2
Abdominal Pain	0.7	2.8	0.0	3.8	2.1
Back Pain	3.0	2.8	0.0	3.8	1.1
Allergic Reaction	2.6	0.9	2.8	1.3	0.0
Asthenia	1.9	2.2	0.0	3.8	0.0
DIGESTIVE SYSTEM					
Constipation	1.8	2.1	0.0	2.5	1.1
Diarrhea	1.5	2.7	0.0	3.8	5.3
Dyspepsia	4.1	2.3	2.8	1.3	2.1
Flatulence	3.3	2.1	2.8	1.3	1.1
RESPIRATORY SYSTEM					
Sinusitis	2.6	2.8	0.0	2.5	6.4
Pharyngitis	1.5	2.5	0.0	1.3	2.1
SKIN AND APPENDAGES					
Rash	0.7	3.9	2.8	3.8	1.1
MUSCULOSKELETAL SYSTEM					
Arthralgia	1.5	2.0	0.0	5.1	0.0
Myalgia	1.1	3.2	5.6	1.3	0.0

Anglo-Scandinavian Cardiac Outcomes Trial (ASCOT): In ASCOT involving 10,305 participants treated with atorvastatin 10 mg daily (n=5,168) or placebo (n=5,137), the safety and tolerability profile of the group treated with atorvastatin was comparable to that of the group treated with placebo during a median of 3.3 years of follow-up. The following adverse events were reported, regardless of causality assessment, in patients treated with atorvastatin in clinical trials. The events in italics occurred in $\geq 2\%$ of patients and the events in plain type occurred in $<2\%$ of patients. **Body as a Whole:** Chest pain, face edema, fever, neck rigidity, malaise, photosensitivity reaction, generalized edema. **Digestive System:** Nausea, gastroenteritis, liver function tests abnormal, colitis, vomiting, gastritis, dry mouth, rectal hemorrhage, esophagitis, eructation, glossitis, mouth ulceration, anorexia, increased appetite, stomatitis, biliary pain, cheilitis, duodenal ulcer, dysphagia, enteritis, melena, gum hemorrhage, stomach ulcer, tenesmus, ulcerative stomatitis, hepatitis, pancreatitis, cholestatic jaundice. **Respiratory System:** Bronchitis, rhinitis, pneumonia, dyspnea, asthma, epistaxis. **Nervous System:** Insomnia, dizziness, paresthesia, somnolence, amnesia, abnormal dreams, libido decreased, emotional lability, incoordination, peripheral neuropathy, torticollis, facial paralysis, hyperkinesia, depression, hypesthesia, hypertonia. **Musculoskeletal System:** Arthritis, leg cramps, bursitis, tenosynovitis, myasthenia, tendinous contracture, myositis. **Skin and Appendages:** Pruritus, contact dermatitis, alopecia, dry skin, sweating, acne, urticaria, eczema, seborrhea, skin ulcer. **Urogenital System:** Urinary tract infection, urinary frequency, cystitis, hematuria, impotence, dysuria, kidney calculus, nocturia, epididymitis, fibrocystic breast, vaginal hemorrhage, albuminuria, breast enlargement, metrorrhagia, nephritis, urinary incontinence, urinary retention, urinary urgency, abnormal ejaculation, uterine hemorrhage. **Special Senses:** Amblyopia, tinnitus, dry eyes, refraction disorder, eye hemorrhage, deafness, glaucoma, parosmia, taste loss, taste perversion. **Cardiovascular System:** Palpitation, vasodilatation, syncope, migraine, postural hypotension, phlebitis, arrhythmia, angina pectoris, hypertension. **Metabolic and Nutritional Disorders:** Peripheral edema, hyperglycemia, creatine phosphokinase increased, gout, weight gain, hypoglycemia. **Hemic and Lymphatic System:** Eosinophilia, anemia, lymphadenopathy, thrombocytopenia, petechia. **Postintroduction Reports with Atorvastatin:** Adverse events associated with atorvastatin therapy reported since market introduction, that are not listed above, regardless of causality assessment, include the following: anaphylaxis, angioneurotic edema, bullous rashes (including erythema multiforme, Stevens-Johnson syndrome, and toxic epidermal necrolysis), and rhabdomyolysis. **Pediatric Patients (ages 10-17 years):** In a 26-week controlled study in boys and postmenarcheal girls (n=140), the safety and tolerability profile of atorvastatin 10 to 20 mg daily was generally similar to that of placebo (see **PRECAUTIONS, Pediatric Use**).

OVERDOSAGE: There is no information on overdosage with CADUET in humans. **Information on Amlodipine:** Single oral doses of amlodipine maleate equivalent to 40 mg amlodipine/kg and 100 mg amlodipine/kg in mice and rats, respectively, caused deaths. Single oral amlodipine maleate doses equivalent to 4 or more mg amlodipine/kg in dogs (11 or more times the maximum recommended clinical dose on a mg/m² basis) caused a marked peripheral vasodilation and hypotension. Overdosage might be expected to cause excessive peripheral vasodilation with marked hypotension and possibly a reflex tachycardia. In humans, experience with intentional overdosage of amlodipine is limited. Reports of intentional overdosage include a patient who ingested 250 mg and was asymptomatic and was not hospitalized; another (120 mg) was hospitalized, underwent gastric lavage and remained normotensive; the third (105 mg) was hospitalized and had hypotension (90/50 mmHg) which normalized following plasma expansion. A patient who took 70 mg amlodipine and an unknown quantity of benzodiazepine in a suicide attempt developed shock which was refractory to treatment and died the following day with abnormally high benzodiazepine plasma concentration. A case of accidental drug overdose has been documented in a 19-month-old male who ingested 30 mg amlodipine (about 2 mg/kg). During the emergency room presentation, vital signs were stable with no evidence of hypotension, but a heart rate of 180 bpm. Ipecac was administered 3.5 hours after ingestion and on subsequent observation (overnight) no sequelae were noted. If massive overdose should occur, active cardiac and respiratory monitoring should be instituted. Frequent blood pressure measurements are essential. Should hypotension occur, cardiovascular support including elevation of the extremities and the judicious administration of fluids should be initiated. If hypotension remains unresponsive to these conservative measures, administration of vasopressors (such as phenylephrine) should be considered with attention to circulating volume and urine output. Intravenous calcium gluconate may help to reverse the effects of calcium entry blockade. As amlodipine is highly protein bound, hemodialysis is not likely to be of benefit. **Information on Atorvastatin:** There is no specific treatment for atorvastatin overdosage. In the event of an overdose, the patient should be treated symptomatically, and supportive measures instituted as required. Due to extensive drug binding to plasma proteins, hemodialysis is not expected to significantly enhance atorvastatin clearance.

*Based on patient weight of 50 kg.

**These events occurred in less than 1% in placebo-controlled trials, but the incidence of these side effects was between 1% and 2% in all multiple dose studies.

Rx only

Manufactured by:
Pfizer Ireland Pharmaceuticals
Dublin, Ireland

© 2004 Pfizer Ireland Pharmaceuticals

Distributed by:



Pfizer Labs

Division of Pfizer Inc., NY, NY 10017

Rev. 1 October 2004

LN233581

© 2005 Pfizer Inc.

All rights reserved.

U.S. Pharmaceuticals

Perfusion CT Proves Useful In Carotid Artery Stenosis

BY PATRICE WENDLING

Chicago Bureau

CHICAGO — Perfusion computed tomography is a useful modality in the detection of regional brain perfusion deficits in patients with severe internal carotid artery stenosis, Dr. Agnieszka Trojanowska during a poster presentation at the annual meeting of the Radiological Society of North America.

CT perfusion imaging revealed that internal carotid artery stenosis in most cases was associated with brain perfusion deficits ipsilaterally to the stenotic site, and that hypoperfusion tended to improve considerably after stent placement, said Dr. Trojanowska, who also has a PhD.

In the study, 74 patients with symptomatic internal carotid artery stenosis of more than 70% were evaluated with CT perfusion imaging, on average, 70 hours

before carotid stent placement and then 3 days and 6 months after stent placement. The protocol included a non-contrast enhanced transaxial CT of the brain with a 5-mm slice and 5-mm slope and dynamic CT perfusion imaging during administration of 50 mL of contrast medium at 4 mL/s with a 5-second delay.

Before stent placement with embolic protection devices, 84% of patients had perfusion deficits ipsilaterally to the stenotic site. Three days after stent placement, 30% of patients had perfusion deficits, and at 6 months, the deficits had diminished to 6%, said Dr. Trojanowska of the Medical University of Lublin (Poland).

A marked elongation of the mean transit time (6.2-6.8 seconds) was noted at the stenotic site, together with decreased values of cerebral blood flow (40-46 mL/100 g per min) and slightly increased cerebral blood volume (3.2 mL/100 g). ■

Fractional Flow Reserve Can Inform Stenting Decisions

SAN FRANCISCO — Looks can be deceiving when evaluating stenoses for treatment with stenting, Dr. John M. Hodgson said at a cardiovascular imaging conference sponsored by the American College of Cardiology.

Not all stenoses detected on angiography are accompanied by ischemia, said Dr. Hodgson of St. Joseph's Hospital, Phoenix. "Two-thirds of the time, when a patient comes to the cath lab we do not have any functional imaging," he said. "We do not know for sure that the patient has ischemia. And then we're left to interpret these fuzzy, two-dimensional angiograms."

But the relatively new technology of measuring fractional flow reserve (FFR) during catheterization could help physicians make better informed decisions about revascularization and stenting.

In FFR, a pressure transducer is sent into the coronary artery, past the anatomic lesion. FFR is the transstenotic pressure gradient across a stenosis, measured at peak blood flow after the administration of a vasodilator (such as adenosine) and indexed for aortic driving pressure.

The result is a direct measurement of the influence of a specific lesion on blood flow. Only when the FFR is 0.75 or less, indicating a functional blockage of at least 25%, is stenting helpful.

The value of FFR was shown in a prospective randomized trial that indicated that not only is it safe to not revascularize stable lesions that don't limit blood flow more than 25%, but also that it provides better 24-month outcomes than does angiography (Circulation 2001;103:2928-34).

—Robert Finn

N-Acetylcysteine May Curb Contrast-Induced Renal Injury

PHILADELPHIA — Giving N-acetylcysteine as an adjunctive agent may reduce the risk of acute renal injury following contrast imaging procedures in high-risk patients, Dr. Venkatesh Jayaraman reported at the annual meeting of the American Society of Nephrology.

Dr. Jayaraman, a nephrology fellow at Lankenau Hospital in Wynnewood, Pa., and associates reviewed the records of 380 patients who underwent coronary angiography in August 2001-January 2004, to evaluate a N-acetylcysteine protocol that was instituted in 2001.

Patients received 600 mg of oral N-acetylcysteine twice daily on the day before and the day of the procedure and were fol-

lowed for 48 hours. By definition, low-risk patients had a serum creatinine level of 1.5 mg/dL or less; high-risk patients were those with more than 1.5 mg/dL.

Among the 318 low-risk patients, there were 8 cases (3%) of contrast-related acute renal failure (ARF). In the 62 high-risk patients, there were 12 cases of ARF (19%). In the low-risk group, 9% of patients received acetylcysteine, compared with 86% of those in the high-risk group.

The investigators concluded that acetylcysteine had a significant effect on the risk of ARF in high-risk, but not low-risk patients. Only about 10% of the high-risk patients who got the drug developed ARF.

—Alicia Ault