

# Cancer Survivors Need Heart Disease Screening

BY MITCHEL L. ZOLER

Philadelphia Bureau

NEW YORK — Heart disease is the leading cause of noncancer mortality in patients who survive an initial bout with Hodgkin's disease, an example of the importance of monitoring for heart disease in cancer survivors.

The relative risk for cardiovascular disease is about two- to sevenfold higher in Hodgkin's disease survivors compared with

age- and gender-matched people with no cancer history, Dr. Ming Hui Chen said at a symposium on cardiovascular disease in cancer patients sponsored by the University of Texas M.D. Anderson Cancer Center.

Similar spikes in cardiovascular disease rates occur in survivors of other cancer types, a link that is mainly attributable to the chemo- or radiotherapy that cancer patients receive. The sequelae usually don't manifest until at least 15 years after cancer treatment. Surveillance for new-onset car-

diovascular disease is challenging because the disease often presents atypically. Chest pain, for example, may be blunted in these patients because their chest nerves were damaged by radiation treatment and so are less sensitive, said Dr. Chen, associate director of the noninvasive cardiac imaging laboratory at the Brigham and Women's Hospital in Boston.

Dr. Chen and her associates have done cardiovascular disease follow-up studies on 182 patients in the Hodgkin's Disease Car-

diac Study. The patients' median age at enrollment was 43, and at entry into the study they were an average of 15 years removed from their initial Hodgkin's disease treatments. A third of the patients received chemotherapy and the remainder had radiation therapy.

During about 4 years of follow-up, cardiovascular diseases were diagnosed in 12 patients (7%), including cases of coronary artery disease, complete heart block, drug-refractory atrial fibrillation, and valvular dysfunction. "For patients who are aged

40-50, the rate [of cardiovascular disease] is quite high," Dr. Chen said at the meeting, also sponsored by the American College of Cardiology and the Society for Geriatric Cardiology.

Current estimates place more than 10

**The relative risk for CVD is about two- to sevenfold higher in Hodgkin's disease survivors. Similar spikes occur in survivors of other cancer types.**

million cancer survivors in the United States, 60% of whom are adults, she said. Breast cancer survivors make up the largest subgroup, but there are also many who had childhood cancers. About 40% of childhood cancer survivors develop severe, disabling complications 30 or more years after their successful cancer treatments.

Guidelines for monitoring cardiovascular disease in childhood cancers survivors were published last February by the Cardiovascular Disease Task Force of the Children's Oncology Group (Pediatrics 2008;121:e387-96). Dr. Chen was a member of the task force.

The guidelines cited three cancer treatments that need regular screening later in life: treatment with an anthracycline, which can potentially cause cardiomyopathy, arrhythmias, or subclinical left-ventricular dysfunction; mediastinal radiation, which can lead to heart failure, cardiomyopathy, pericarditis, pericardial fibrosis, valvular disease, myocardial infarction, arrhythmia, or atherosclerotic heart disease; and neck radiation of 40 Gy or more, which can lead to carotid and subclavian artery disease.

Patients who undergo any of these treatments should be followed up with an annual history and physical, the guidelines said. In addition, patients who received an anthracycline or mediastinal radiation should receive a baseline echocardiography examination (or multiple gated-acquisition scan if treated with an anthracycline) followed by periodic follow-up echo examinations with the frequency dictated by the patient's age when treated and the cumulative dose received. These patients should also have a baseline electrocardiogram to evaluate their QT interval. Following mediastinal radiation, patients should also be screened with a fasting glucose and lipid profile every 3-5 years. Following neck radiation, patients should undergo Doppler ultrasound examinations of their carotid and subclavian arteries as clinically needed. ■

## OMNARIS™ (ciclesonide)

Nasal Spray, 50 mcg

For intranasal use only

Rx only

**BRIEF SUMMARY:** Please see package insert for full prescribing information.

### INDICATIONS AND USAGE

#### Seasonal Allergic Rhinitis

OMNARIS Nasal Spray is indicated for the treatment of nasal symptoms associated with seasonal allergic rhinitis in adults and children 6 years of age and older.

#### Perennial Allergic Rhinitis

OMNARIS Nasal Spray is indicated for the treatment of nasal symptoms associated with perennial allergic rhinitis in adults and adolescents 12 years of age and older.

### CONTRAINDICATIONS

OMNARIS Nasal Spray is contraindicated in patients with a hypersensitivity to any of its ingredients.

### WARNINGS

The replacement of a systemic corticosteroid with a topical corticosteroid can be accompanied by signs of adrenal insufficiency. In addition, some patients may experience symptoms of corticosteroid withdrawal, e.g., joint and/or muscular pain, lassitude, and depression. Patients previously treated for prolonged periods with systemic corticosteroids and transferred to topical corticosteroids should be carefully monitored for acute adrenal insufficiency in response to stress. In those patients who have asthma or other clinical conditions requiring long-term systemic corticosteroid treatment, rapid decreases in systemic corticosteroid dosages may cause a severe exacerbation of their symptoms.

Patients who are using drugs that suppress the immune system are more susceptible to infections than healthy individuals. Chickenpox and measles, for example, can have a more serious or even fatal course in children or adults using corticosteroids. In children or adults who have not had these diseases or been properly immunized, particular care should be taken to avoid exposure. How the dose, route, and duration of corticosteroid administration affect the risk of developing a disseminated infection is not known. The contribution of the underlying disease and/or prior corticosteroid treatment to the risk is also not known. If exposed to chickenpox, prophylaxis with varicella zoster immune globulin (VZIG) may be indicated. If exposed to measles, prophylaxis with pooled intramuscular immunoglobulin (IG) may be indicated. (See the respective package inserts for complete VZIG and IG prescribing information.) If chickenpox develops, treatment with antiviral agents may be considered.

### PRECAUTIONS

#### General

Intranasal corticosteroids may cause a reduction in growth velocity when administered to pediatric patients (see PRECAUTIONS: Pediatric Use). Rarely, immediate hypersensitivity reactions or contact dermatitis may occur after the administration of intranasal corticosteroids. Patients with a known hypersensitivity reaction to other corticosteroid preparations should use caution when using ciclesonide nasal spray since cross reactivity to other corticosteroids including ciclesonide may also occur.

Because of the inhibitory effect of corticosteroids on wound healing, patients who have experienced recent nasal septal ulcers, nasal surgery, or nasal trauma should not use a nasal corticosteroid until healing has occurred. In clinical studies with OMNARIS Nasal Spray, the development of localized infections of the nose and pharynx with *Candida albicans* has rarely occurred. When such an infection develops, it may require treatment with appropriate local therapy and discontinuation of OMNARIS Nasal Spray. Therefore, patients using OMNARIS Nasal Spray over several months or longer should be examined periodically for evidence of *Candida* infection or other signs of adverse effects on the nasal mucosa. Intranasal corticosteroids should be used with caution, if at all, in patients with active or quiescent tuberculosis infections of the respiratory tract; or in patients with untreated local or systemic fungal or bacterial infections; systemic viral or parasitic infections; or ocular herpes simplex.

If recommended doses of intranasal corticosteroids are exceeded or if individuals are particularly sensitive or predisposed by virtue of recent systemic steroid therapy, symptoms of hypercorticism may occur, including very rare cases of menstrual irregularities, acneiform lesions, and cushingoid features. If such changes occur, topical corticosteroids should be discontinued slowly, consistent with accepted procedures for discontinuing oral steroid therapy.

The risk of glaucoma was evaluated by assessments of intraocular pressure in 3 studies including 943 patients. Of these, 390 adolescents or adults were treated for up to 52 weeks and 186 children ages 2 to 11 received treatment with OMNARIS Nasal Spray 200 mcg daily for up to 12 weeks. In these trials, no significant differences in intraocular pressure changes were observed between OMNARIS Nasal Spray 200 mcg and placebo-treated patients. Additionally, no significant differences between OMNARIS Nasal Spray 200 mcg and placebo-treated patients were noted during the 52-week study of adults and adolescent patients in whom thorough ophthalmologic assessments were performed including evaluation of cataract formation using slit lamp examinations. Rare instances of wheezing, nasal septum perforation, cataracts, glaucoma, and increased intraocular pressure have been reported following the intranasal application of corticosteroids. Close follow-up is warranted in patients with a change in vision and with a history of glaucoma and/or cataracts.

#### Information for Patients

Patients being treated with OMNARIS Nasal Spray should receive the following information and instructions. This information is intended to aid them in the safe and effective use of this medication. It is not a disclosure of all possible adverse or intended effects.

Patients who are on immunosuppressive doses of corticosteroids should be warned to avoid exposure to chickenpox or measles, and if exposed, to obtain medical advice. Patients should use OMNARIS Nasal Spray at regular intervals since its effectiveness depends on its regular use (see DOSAGE AND ADMINISTRATION).

In clinical trials, the onset of effect was seen within 24 to 48 hours with further symptomatic improvement observed over 1 to 2 weeks in seasonal allergic rhinitis and 5 weeks in perennial allergic rhinitis. Initial assessment of response should be made during this timeframe and periodically until the patients symptoms are stabilized.

The patient should take the medication as directed and should not exceed the prescribed dosage. The patient should contact the physician if symptoms do not improve by a reasonable time or if the condition worsens. For the proper use of this unit and to attain maximum improvement, the patients should read and follow the accompanying patient instructions carefully. Spraying OMNARIS Nasal Spray directly into the eyes or onto the nasal septum should be avoided. It is important that the bottle is gently shaken prior to use to ensure that a consistent amount is dispensed per actuation. The bottle should be discarded after 120 actuations following initial priming or after 4 months after the bottle is removed from the foil pouch, whichever occurs first.

#### Drug Interactions

Based on *in vitro* studies in human liver microsomes, des-ciclesonide appears to have no inhibitory or induction potential on the metabolism of other drugs metabolized by CYP 450 enzymes. The inhibitory potential of ciclesonide on CYP450 isoenzymes has not been studied. *In vitro* studies demonstrated that the plasma protein binding of des-ciclesonide was not affected by warfarin or salicylic acid, indicating no potential for protein binding-based drug interactions.

In a drug interaction study, co-administration of orally inhaled ciclesonide and oral erythromycin, an inhibitor of cytochrome P450 3A4, had no effect on the pharmacokinetics of either des-ciclesonide or erythromycin. In another drug interaction study, co-administration of orally inhaled ciclesonide and oral ketoconazole, a potent inhibitor of cytochrome P450 3A4, increased the exposure (AUC) of des-ciclesonide by approximately 3.6-fold at steady state, while levels of ciclesonide remained unchanged. Therefore, ketoconazole should be administered with caution with intranasal ciclesonide.

#### Carcinogenesis, Mutagenesis, Impairment of Fertility

Ciclesonide demonstrated no carcinogenic potential in a study of oral doses up to 900 mcg/kg (approximately 20 to 10 times the maximum human daily intranasal dose in adults and children, respectively, based on mcg/m<sup>2</sup>) in mice for 104 weeks and in a study of inhalation doses up to 193 mcg/kg (approximately 8 and 5 times the maximum human daily intranasal dose in adults and children, respectively, based on mcg/m<sup>2</sup>) in rats for 104 weeks. Ciclesonide was not mutagenic in an Ames test or in a forward mutation assay and was not clastogenic in a human lymphocyte assay or in an *in vitro* micronucleus test. However, ciclesonide was clastogenic in the *in vivo* mouse micronucleus test. The concurrent reference corticosteroid (dexamethasone) in this study showed similar findings. No evidence of impairment of fertility was observed in a reproductive study conducted in male and female rats both dosed orally up to

900 mcg/kg/day (approximately 35 times the maximum human daily intranasal dose in adults based on mcg/m<sup>2</sup>).

#### Pregnancy: Teratogenic Effects

##### Pregnancy Category C

Oral administration of ciclesonide in rats up to 900 mcg/kg (approximately 35 times the maximum human daily intranasal dose in adults based on mcg/m<sup>2</sup>) produced no teratogenicity or other fetal effects. However, subcutaneous administration of ciclesonide in rabbits at 5 mcg/kg (less than the maximum human daily intranasal dose in adults based on mcg/m<sup>2</sup>) or greater produced fetal toxicity. This included fetal loss, reduced fetal weight, cleft palate, skeletal abnormalities including incomplete ossifications, and skin effects. No toxicity was observed at 1 mcg/kg (less than the maximum human daily intranasal dose based on mcg/m<sup>2</sup>).

There are no adequate and well-controlled studies in pregnant women. OMNARIS Nasal Spray, like other corticosteroids, should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Experience with oral corticosteroids since their introduction in pharmacologic, as opposed to physiologic, doses suggests that rodents are more prone to teratogenic effects from corticosteroids than humans. In addition, because there is a natural increase in corticosteroid production during pregnancy, most women will require a lower exogenous corticosteroid dose and many will not need corticosteroid treatment during pregnancy.

#### Nonteratogenic Effects

Hypoadrenalism may occur in infants born of mothers receiving corticosteroids during pregnancy. Such infants should be carefully monitored.

#### Nursing Mothers

It is not known if ciclesonide is excreted in human milk. However, other corticosteroids are excreted in human milk. In a study with lactating rats, minimal but detectable levels of ciclesonide were recovered in milk. Caution should be used when OMNARIS Nasal Spray is administered to nursing women.

#### Pediatric Use

The safety and effectiveness for seasonal and perennial allergic rhinitis in children 12 years of age and older have been established. The efficacy of OMNARIS Nasal Spray in patients 6 to 11 years of age for treatment of the symptoms of seasonal allergic rhinitis is supported by evidence from four adequate and well-controlled studies in adults and adolescents 12 years of age and older with seasonal and perennial allergic rhinitis, and one study in patients 6 to 11 years of age with seasonal allergic rhinitis. The efficacy of OMNARIS Nasal Spray for the treatment of the symptoms of perennial allergic rhinitis in patients 6 to 11 years of age has not been established (see CLINICAL TRIALS: Pediatric Patients Aged 6 to 11 Years). The efficacy of OMNARIS Nasal Spray in children 2 to 5 years of age has not been established. The safety of OMNARIS Nasal Spray in children 2 to 11 years of age was evaluated in 4 controlled clinical studies of 2 to 12 weeks duration (see CLINICAL PHARMACOLOGY: Pharmacodynamics, CLINICAL TRIALS, ADVERSE REACTIONS: Pediatric Patients).

Clinical studies in children less than two years of age have not been conducted. Studies in children under 2 years of age are waived because of local and systemic safety concerns.

Controlled clinical studies have shown that intranasal corticosteroids may cause a reduction in growth velocity in pediatric patients. This effect has been observed in the absence of laboratory evidence of hypothalamic-pituitary-adrenal (HPA)-axis suppression, suggesting that growth velocity is a more sensitive indicator of systemic corticosteroid exposure in pediatric patients than some commonly used tests of HPA-axis function. The long-term effects of this reduction in growth velocity associated with intranasal corticosteroids, including the impact on final adult height, are unknown. The potential for "catch-up" growth following discontinuation of treatment with intranasal corticosteroids has not been adequately studied. The growth of pediatric patients receiving intranasal corticosteroids, including OMNARIS Nasal Spray, should be monitored routinely (e.g., via stadiometry). The potential growth effects of prolonged treatment should be weighed against clinical benefits obtained and the availability of safe and effective noncorticosteroid treatment alternatives. To minimize the systemic effects of intranasal corticosteroids, each patient should be titrated to the lowest dose that effectively controls his/her symptoms.

#### Geriatric Use

Clinical studies of OMNARIS Nasal Spray did not include sufficient numbers of subjects age 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

#### ADVERSE REACTIONS

##### Adult and Adolescent Patients Aged 12 Years and Older:


In controlled clinical studies conducted in the US and Canada, a total of 1524 patients ages 12 years and older received treatment with ciclesonide administered intranasally. The overall incidence of adverse events for patients treated with OMNARIS Nasal Spray was comparable to that in patients treated with placebo. Adverse events did not differ appreciably based on age, gender, or race. Approximately 2% of patients treated with OMNARIS Nasal Spray 200 mcg in clinical trials discontinued because of adverse events; this rate was similar for patients treated with placebo. Adverse events, irrespective of drug relationship, that occurred with an incidence of 2% or greater and more frequently with OMNARIS Nasal Spray 200 mcg (N = 546) than with placebo (N = 544) in clinical trials of 2 to 6 weeks in duration included headache (6.0% vs 4.6%), epistaxis (4.9% vs 2.9%), nasopharyngitis (3.7% vs 3.3), and ear pain (2.2% vs 0.6%). In a 52-week long-term safety trial that included 663 adults and adolescent patients (441 treated with ciclesonide: 227 males and 436 females) with perennial allergic rhinitis, the adverse event profile over the treatment period was similar to the adverse event profile in trials of shorter duration. Adverse events considered likely or definitely related to OMNARIS Nasal Spray that were reported at an incidence of 1% or greater of patients and more commonly in OMNARIS Nasal Spray versus placebo were epistaxis, nasal discomfort, and headache. No patient experienced a nasal septal perforation or nasal ulcer during long-term use of OMNARIS Nasal Spray. While primarily designed to assess the long-term safety of OMNARIS Nasal Spray 200 mcg once daily, this 52-week trial demonstrated greater decreases in total nasal symptom scores with OMNARIS Nasal Spray versus placebo treated patients over the entire treatment period.

##### Pediatric Patients Aged 6 to 11 Years:

Two controlled clinical studies 2 and 12 weeks in duration were conducted in the US and Canada and included a total of 1282 patients with allergic rhinitis ages 6 to 11 years, of which 913 were treated with OMNARIS (ciclesonide) Nasal Spray 200 mcg, 100 mcg, or 25 mcg daily. The overall incidence of adverse events for patients treated with OMNARIS Nasal Spray was comparable to that in patients treated with placebo. Adverse events did not differ appreciably based on age, gender, or race. In clinical trials, 1.6% and 2.7% of patients treated with OMNARIS Nasal Spray 200 mcg or 100 mcg, respectively, discontinued because of adverse events; these rates were lower than the rate in patients treated with placebo (2.8%). Adverse events, irrespective of drug relationship, that occurred with an incidence of 3% or greater and more frequently with OMNARIS Nasal Spray 200 mcg (N = 380) than with placebo (N = 369) included headache (6.6% vs 5.7%), nasopharyngitis (6.6% vs 5.4%), and pharyngolaryngeal pain (3.4% vs 3.3%).

##### Pediatric Patients Aged 2 to 5 Years:

Two controlled clinical studies 6 and 12 weeks in duration were conducted in the US and included a total of 258 patients 2 to 5 years of age with perennial allergic rhinitis, of which 183 were treated with OMNARIS Nasal Spray 200 mcg, 100 mcg or 25 mcg daily. The distribution of adverse events was similar to that seen in the 6 to 11 year old children.



Manufactured for:  
Sepracor Inc.  
Marlborough, MA 01752 USA  
Made in Germany  
November 2007

© 2008 Sepracor Inc. Marlborough, MA 01752 All rights reserved. 4/08 OMN135-08



and SEPRACOR are registered trademarks of Sepracor Inc.

OMNARIS is a trademark of Nycomed GmbH, used with permission.