

# Eating Flaxseed May Slow Prostate Cancer Growth

BY KERRI WACHTER  
Senior Writer

CHICAGO — Adding flaxseed to the diets of men with prostate cancer appears to slow tumor growth, according to data presented at the annual meeting of the American Society of Clinical Oncology.

In a study of 161 men with prostate cancer, those who consumed 30 g of ground flaxseed daily had significantly lower levels of the tumor proliferation marker

MIB-1 than did men on their usual diet.

"Prostate cancer proliferation rates were significantly lower in men assigned to flaxseed supplementation," said Wendy Demark-Wahnefried, Ph.D., a researcher in the school of nursing at Duke University, Durham, N.C.

Flaxseed is a rich source of lignans, which have been shown to affect androgen metabolism and have antimetabolic, antiangiogenic, antioxidant, and estrogenic effects. Flaxseed also is a rich source of

omega-3 fatty acids, which play a role in cell membrane function, inhibit protein kinase activity, and increase natural killer-cell activity. Previous studies have suggested, too, that a low-fat diet might help prevent prostate cancer.

The study included men with pathologically confirmed prostate cancer who were electing prostatectomy as their primary treatment. Their mean age was 59 years, and 70% were white. They had to be at least 21 days from scheduled surgery

and not on neoadjuvant therapy. The men could not have started any new supplements (excluding multivitamins) during or within 3 months of the study period. Also, they could not use antibiotics during or 7 days prior to the study, because normal gut flora is necessary to process lignans.

The men were randomized to supplementation with flaxseed (40 patients), a low-fat diet (40), flaxseed plus a low-fat diet (40), or their usual diet (41). Men on their usual diet served as controls. Men receiving flaxseed supplementation (donated by Enreco Inc.) consumed 30 g of ground flaxseed daily. Men on a low-fat diet were limited to receiving less than 20% of daily calories from fat.

At baseline and immediately prior to surgery, blood, urine, and seminal fluid were collected, and the men completed questionnaires. Following surgery, pro-



**Brief Summary:** For complete details, please see full Prescribing Information.

**INDICATIONS AND USAGE:** BYETTA is indicated as adjunctive therapy to improve glycemic control in patients with type 2 diabetes mellitus who are taking metformin, a sulfonylurea, a thiazolidinedione, a combination of metformin and a sulfonylurea, or a combination of metformin and a thiazolidinedione, but have not achieved adequate glycemic control.

**CONTRAINDICATIONS:** BYETTA is contraindicated in patients with known hypersensitivity to exenatide or to any of the product components.

**PRECAUTIONS:** General—BYETTA is not a substitute for insulin in insulin-requiring patients. BYETTA should not be used in patients with type 1 diabetes or for the treatment of diabetic ketoacidosis.

Patients may develop anti-exenatide antibodies following treatment with BYETTA, consistent with the potentially immunogenic properties of protein and peptide pharmaceuticals. Patients receiving BYETTA should be observed for signs and symptoms of hypersensitivity reactions. In a small proportion of patients, the formation of anti-exenatide antibodies at high titers could result in failure to achieve adequate improvement in glycemic control.

The concurrent use of BYETTA with insulin, D-phenylalanine derivatives, meglitinides, or alpha-glucosidase inhibitors has not been studied.

BYETTA is not recommended for use in patients with end-stage renal disease or severe renal impairment (creatinine clearance <30 mL/min; see Pharmacokinetics, Special Populations). In patients with end-stage renal disease receiving dialysis, single doses of BYETTA 5 mcg were not well tolerated due to gastrointestinal side effects.

BYETTA has not been studied in patients with severe gastrointestinal disease, including gastroparesis. Its use is commonly associated with gastrointestinal adverse effects, including nausea, vomiting, and diarrhea. Therefore, the use of BYETTA is not recommended in patients with severe gastrointestinal disease. The development of severe abdominal pain in a patient treated with BYETTA should be investigated because it may be a warning sign of a serious condition.

**Hypoglycemia**—In the 30-week controlled clinical trials with BYETTA, a hypoglycemia episode was recorded as an adverse event if the patient reported symptoms associated with hypoglycemia with an accompanying blood glucose <60 mg/dL or if symptoms were reported without an accompanying blood glucose measurement. When BYETTA was used in combination with metformin, no increase in the incidence of hypoglycemia was observed. In contrast, when BYETTA was used in combination with a sulfonylurea, the incidence of hypoglycemia was increased over that of placebo in combination with a sulfonylurea. Therefore, patients receiving BYETTA in combination with a sulfonylurea may have an increased risk of hypoglycemia (Table 1).

Table 1: Incidence (%) of Hypoglycemia\* by Concomitant Antidiabetic Therapy

	BYETTA			BYETTA			BYETTA		
	Placebo BID	5 mcg BID	10 mcg BID	Placebo BID	5 mcg BID	10 mcg BID	Placebo BID	5 mcg BID	10 mcg BID
	With Metformin			With a Sulfonylurea			With MET/SFU		
N	113	110	113	123	125	129	247	245	241
Hypoglycemia	5.3%	4.5%	5.3%	3.3%	14.4%	35.7%	12.6%	19.2%	27.8%

\* In three 30-week placebo-controlled clinical trials. BYETTA and placebo were administered before the morning and evening meals. Abbreviations: BID, twice daily; MET/SFU, metformin and a sulfonylurea.

Most episodes of hypoglycemia were mild to moderate in intensity, and all resolved with oral administration of carbohydrate. To reduce the risk of hypoglycemia associated with the use of a sulfonylurea, reduction in the dose of sulfonylurea may be considered (see DOSAGE AND ADMINISTRATION). When used as add-on to a thiazolidinedione, with or without metformin, the incidence of symptomatic mild to moderate hypoglycemia with BYETTA was 11% compared to 7% with placebo.

BYETTA did not alter the counter-regulatory hormone responses to insulin-induced hypoglycemia in a randomized, double-blind, controlled study in healthy subjects.

**Information for Patients**—Patients should be informed of the potential risks of BYETTA. Patients should also be fully informed about self-management practices, including the importance of proper storage of BYETTA, injection technique, timing of dosage of BYETTA as well as concomitant oral drugs, adherence to meal planning, regular physical activity, periodic blood glucose monitoring and HbA<sub>1c</sub> testing, recognition and management of hypoglycemia and hyperglycemia, and assessment for diabetes complications.

Patients should be advised to inform their physicians if they are pregnant or intend to become pregnant.

The risk of hypoglycemia is increased when BYETTA is used in combination with an agent that induces hypoglycemia, such as a sulfonylurea (see PRECAUTIONS, Hypoglycemia).

Patients should be advised that treatment with BYETTA may result in a reduction in appetite, food intake, and/or body weight, and that there is no need to modify the dosing regimen due to such effects. Treatment with BYETTA may also result in nausea (see ADVERSE REACTIONS).

**Drug Interactions**—The effect of BYETTA to slow gastric emptying may reduce the extent and rate of absorption of orally administered drugs. BYETTA should be used with caution in patients receiving oral medications that require rapid gastrointestinal absorption. For oral medications that are dependent on threshold concentrations for efficacy, such as contraceptives and antibiotics, patients should be advised to take those drugs at least 1 h before BYETTA injection. If such drugs are to be administered with food, patients should be advised to take them with a meal or snack when BYETTA is not administered. The effect of BYETTA on the absorption and effectiveness of oral contraceptives has not been characterized.

**Warfarin:** Since market introduction there have been some spontaneously reported cases of increased INR with concomitant use of warfarin and BYETTA, sometimes associated with bleeding.

**Carcinogenesis, Mutagenesis, Impairment of Fertility**—A 104-week carcinogenicity study was conducted in male and female rats and benign thyroid C-cell adenomas were observed in female rats at all exenatide doses. The incidences in female rats were 8% and 5% in the two control groups and 14%, 11%, and 23% in the low-, medium-, and high-dose groups with systemic exposures of 5, 22, and 130 times, respectively, the human exposure resulting from the maximum recommended dose of 20 mcg/day.

In a 104-week carcinogenicity study in mice, no evidence of tumors was observed at doses up to 250 mcg/kg/day, a systemic exposure up to 95 times the human exposure resulting from the maximum recommended dose of 20 mcg/day.

Exenatide was not mutagenic or clastogenic, with or without metabolic activation, in the Ames bacterial mutagenicity assay or chromosomal aberration assay in Chinese hamster ovary cells.

**Pregnancy—Pregnancy Category C**—Exenatide has been shown to cause reduced fetal and neonatal growth, and skeletal effects in mice at systemic exposures 3 times the human exposure resulting from the maximum recommended dose of 20 mcg/day. Exenatide has been shown to cause skeletal effects in rabbits at systemic exposures 12 times the human exposure resulting from the maximum recommended dose of 20 mcg/day. There are no adequate and well-controlled studies in pregnant women. BYETTA should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

In pregnant mice an increased number of neonatal deaths were observed on postpartum days 2-4 in dams given 6 mcg/kg/day, a systemic exposure 3 times the human exposure resulting from the maximum recommended dose of 20 mcg/day.

**Nursing Mothers**—It is not known whether exenatide is excreted in human milk. Caution should be exercised when BYETTA is administered to a nursing woman.

**Pediatric Use**—Safety and effectiveness of BYETTA have not been established in pediatric patients.

**Geriatric Use**—BYETTA was studied in 282 patients 65 years of age or older and in 16 patients 75 years of age or older. No differences in safety or effectiveness were observed between these patients and younger patients.

**ADVERSE REACTIONS: Use with metformin and/or a sulfonylurea**—In the three 30-week controlled trials of BYETTA add-on to metformin and/or sulfonylurea, adverse events with an incidence  $\geq$  5% (excluding hypoglycemia; see Table 1) that occurred more frequently in patients treated with BYETTA (N = 963) vs placebo (N = 483) were: nausea (44% vs 18%), vomiting (13% vs 4%), diarrhea (13% vs 6%), feeling jittery (9% vs 4%), dizziness (9% vs 6%), headache (9% vs 6%), and dyspepsia (6% vs 3%).

The adverse events associated with BYETTA generally were mild to moderate in intensity. The most frequently reported adverse event, mild to moderate nausea, occurred in a dose-dependent fashion. With continued therapy, the frequency and severity decreased over time in most of the patients who initially experienced nausea. Adverse events reported in  $\sim$  1.0 to  $<$  5.0% of patients receiving BYETTA and reported more frequently than with placebo included asthenia (mostly reported as weakness), decreased appetite, gastroesophageal reflux disease, and hyperhidrosis. Patients in the extension studies at 52 weeks experienced similar types of adverse events observed in the 30-week controlled trials.

The incidence of withdrawal due to adverse events was 7% for BYETTA-treated patients and 3% for placebo-treated patients. The most common adverse events leading to withdrawal for BYETTA-treated patients were nausea (3% of patients) and vomiting (1%). For placebo-treated patients,  $<$  1% withdrew due to nausea and 0% due to vomiting.

**Use with a thiazolidinedione**—In the 16-week placebo-controlled study of BYETTA add-on to a thiazolidinedione, with or without metformin, the incidence and type of other adverse events observed were similar to those seen in the 30-week controlled clinical trials with metformin and/or a sulfonylurea. No serious adverse events were reported in the placebo arm. Two serious adverse events, namely chest pain (leading to withdrawal) and chronic hypersensitivity pneumonitis, were reported in the BYETTA arm.

The incidence of withdrawal due to adverse events was 16% (19/121) for BYETTA-treated patients and 2% (2/112) for placebo-treated patients. The most common adverse events leading to withdrawal for BYETTA-treated patients were nausea (9%) and vomiting (5%). For placebo-treated patients,  $<$  1% withdrew due to nausea. Chills (n = 4) and injection-site reactions (n = 2) occurred only in BYETTA-treated patients. The two patients who reported an injection-site reaction had high titers of anti-exenatide antibody.

**Spontaneous Data**—Since market introduction of BYETTA, the following additional adverse reactions have been reported. Because these events are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. **General:** injection-site reactions; dysgeusia; somnolence, INR increased with concomitant warfarin use (some reports associated with bleeding). **Allergy/Hypersensitivity:** generalized pruritus and/or urticaria, macular or papular rash, angioedema; rare reports of anaphylactic reaction. **Gastrointestinal:** nausea, vomiting, and/or diarrhea resulting in dehydration with some reports associated with increased serum creatinine/acute renal failure that may be reversible if treated appropriately; abdominal distension, abdominal pain, eructation, constipation, flatulence, acute pancreatitis.

**Immunogenicity**—Consistent with the potentially immunogenic properties of protein and peptide pharmaceuticals, patients may develop anti-exenatide antibodies following treatment with BYETTA.

**OVERDOSAGE:** Effects of an overdose include severe nausea, severe vomiting, and rapidly declining blood glucose concentrations. In the event of overdose, appropriate supportive treatment should be initiated according to the patient's clinical signs and symptoms.

**DOSAGE AND ADMINISTRATION:** BYETTA therapy should be initiated at 5 mcg per dose administered twice daily at any time within the 60-minute period before the morning and evening meals (or before the two main meals of the day, approximately 6 hours or more apart). BYETTA should not be administered after a meal. Based on clinical response, the dose of BYETTA can be increased to 10 mcg twice daily after 1 month of therapy. Each dose should be administered as a SC injection in the thigh, abdomen, or upper arm.

#### Rx ONLY

Manufactured by Amylin Pharmaceuticals, Inc., San Diego, CA 92121  
Marketed by Amylin Pharmaceuticals, Inc. and Eli Lilly and Company  
1-800-868-1190  
Literature Revised February 2007

BYETTA is a registered trademark of Amylin Pharmaceuticals, Inc.  
© 2007 Amylin Pharmaceuticals, Inc. All rights reserved.

www.BYETTA.com  
02-05-1649-G

822001-GG



**Flaxseed might reduce cancer proliferation by altering androgen metabolism.**

**DR. DEMARK-WAHNEFRIED**

static tissue was analyzed. Men were stratified by race and combined Gleason score (classified as either above or below 7) so that the groups had similar representation of aggressive disease. Overall, 68% of men had a Gleason score of less than 7.

Men in all four groups followed their protocols for roughly 30 days. Adherence (confirmed by biomarkers) was high in the three intervention groups: almost 7 days per week. The attrition rate of 9% was due mainly to men deciding against surgery, or advancement of their surgery date.

Men in the flaxseed-only group and those in the combination flaxseed/low-fat diet group had significantly lower levels of MIB-1, indicating lower rates of tumor proliferation, expressed as the ratio of cancer cells that are actively dividing to those that are not dividing: 1.66 in the flaxseed group and 1.5 in the combination group, compared with 2.56 in the low-fat diet group and 3.23 in the control group. Other biomarkers that are associated with prostate cancer, including apoptosis and androgen metabolism, did differ significantly.

"Unfortunately, we did not see what we'd been able to find in our pilot studies, which was a reduction in testosterone levels and PSA [prostate-specific antigen] in our flaxseed groups," Dr. Demark-Wahnefried said. "In our control group, the PSA and androgen levels really bottomed out. So there were no differences detected."

The researchers had speculated that flaxseed might reduce cancer proliferation by altering androgen metabolism, she said. They hope to identify the mechanism in future studies.

There were no differences between the groups in terms of side effects. ■