

## Question the Right People For Bipolar Dx

BY SHERRY BOSCHERT  
San Francisco Bureau

SAN FRANCISCO — Diagnosing bipolar disorder requires not just asking the right questions but the right people, Michael J. Gitlin, M.D., said.

More than a third of 600 bipolar disorder patients sought help within a year of becoming symptomatic, but 69% were misdiagnosed (usually as having major depression). In addition, patients saw a mean of four physicians before being correctly diagnosed, a 2003 survey found.

"My quarrel with these data that get published over and over is that the implication is that the physicians" erred, said Dr. Gitlin, professor of clinical psychiatry at the University of California, Los Angeles. "I think it's more complex than that."

Even physicians who ask the right questions often fail to get the whole story from bipolar disorder patients. "You need corroborative evidence" obtained by getting the patient's permission to talk with significant others—a spouse, boyfriend or girlfriend, sibling, or parents, he said.

The same survey of 600 bipolar disorder patients found that they reported manic symptoms far less often than they reported depressive symptoms to physicians when seeking help (*J. Clin. Psychiatry* 2003;64:161-74).

That's not surprising, he noted, because depression feels bad and brings people into treatment. Patients reported manic symptoms other than erratic sleep only 43% of the time or less. Only 37% of patients, for example, told physicians of feeling elated at times.

"That is much more the issue than the idea that we're not probing correctly," he said.

Bipolar disorder patients may not recognize manic or hypomanic symptoms as abnormal, or remember having them. When they're depressed, they're dominated by depressive thoughts and symptoms, and when they're hypomanic they may not remember the depressive symptoms.

Ask significant others as well as the patient about any hypomanic symptoms immediately before or after depressed periods. The highs and lows of bipolar disorder are temporarily related, usually with mania preceding depression, but less commonly in reverse order, Dr. Gitlin said.

Don't just focus on mood, but ask about overactivity, which may be a core feature of hypomania. "The data are not entirely convincing for this, but I think it's something we really ought to be thinking about," he said.

Ask about a family history of mood disorders, and ask whether the patient or family members have ever shown signs of antidepressant-induced hypomania. If you decide to start an antidepressant, educate the patient and significant others to watch for signs of induced hypomania. ■

## Subthreshold Bipolar Disorder Is Prevalent

BY MITCHEL L. ZOLER  
Philadelphia Bureau

PITTSBURGH — Subthreshold bipolar disorder is highly prevalent and disabling, according to a nationwide survey of more than 9,000 Americans sponsored by the National Institute of Mental Health.

About 2% of the U.S. adult population has subthreshold bipolar disorder, on top of the 2% who have symptoms that meet DSM-IV criteria for bipolar I and II dis-


order, Ronald C. Kessler, Ph.D., said at the Sixth International Conference on Bipolar Disorder. These are "staggeringly large numbers," he said.

The survey results also documented that people who reported having subthreshold bipolar disorder, defined as having episodes with two or more symptoms of bipolar disorder for 4 or more days at a time, had an average of 43 days a year when they were totally unable to work or to perform other normal, daily

activities. This level of impairment was very similar to what was reported by people who met the definition for classic bipolar I or II disorder.

"We need to figure out how we can deliver effective treatment to people" with subthreshold bipolar disorder, said Dr. Kessler, a professor in the department of health care policy at Harvard Medical School in Boston. "The subthreshold people sure have something that's much more impairing than most things we call

INTRODUCING  
A **NEW** ONCE-DAILY D-GREE OF RELIEF



# POWER UNIMPAIRED

ALL THE POWER OF ALLEGRA® 180 MG  
**PLUS** 24 HOURS OF CONGESTION RELIEF

**NEW!** 24 Hour  
*allegra-D*®

fexofenadine HCl 180 mg/pseudoephedrine HCl 240 mg  
Extended-Release Tablets