

# Frovatriptan Relieves Pure Menstrual Migraines

BY HEIDI SPLETE  
Senior Writer

WASHINGTON — Women who took 2.5 mg of frovatriptan either once or twice daily for 6 days at the time of menstruation had significantly fewer—and less severe—menstrual migraines, compared with women who took a placebo, reported Dr. Marie Pinizzotto and her colleagues at Endo Pharmaceuticals.

The women on either regimen of frova-

triptan also reported significantly fewer headaches in general and less functional impairment compared with the placebo group. Data from the randomized, double-blind, three-way crossover study were presented in a poster at the annual meeting of the American College of Obstetricians and Gynecologists. The study was sponsored by Vernalis Development Ltd., and Endo Pharmaceuticals Inc.

Frovatriptan has been approved by the Food and Drug Administration for the acute

treatment of migraines, both with and without aura, in adults, but it has not been approved for the prophylactic prevention of migraines. The manufacturers are seeking an additional indication for the prophylactic treatment of menstrual migraines.

The patients were randomized to receive each of the two treatment regimens or a placebo over the course of three different 6-day periods from 2 days before to 4 days after the onset of menstruation.

The incidence of pure menstrual mi-

graines, defined as migraines that occurred during the time period from 2 days before to 3 days after the onset of menstruation, was significantly lower in both frovatriptan groups, compared with placebo. These distinctive headaches occurred in 38% of the twice-daily frovatriptan group, compared with 51% of the once-daily group and 67% of the placebo group.

The intent-to-treat analysis included 179 women aged 18 years and older with at least a 1-year history of menstrually-related migraines. The mean age was 37 years, and 82% were white. On average, the study participants had a history of migraines greater than 10 years, and the average number of migraine attacks was one per month during the year prior to the study.

Moderate to severe headaches were reported by 25%, 32%, and 46% of women in the twice-daily frovatriptan, once-daily frovatriptan, and placebo groups. The incidence of functional impairment was 14%, 24%, and 35%, respectively.

Adverse events included headache, nausea, dizziness, and nasopharyngitis, and the incidence of these events was similar between the two groups, with the exception of upper respiratory tract infections, which were significantly more common in the patients treated with twice-daily frovatriptan. ■

## Perinatal Risks Increase With Maternal Weight

WASHINGTON — Either too much or too little weight gain during pregnancy could increase the risk of neonatal intensive care unit admission and peripartum complications, according to data presented at the annual meeting of the American College of Obstetricians and Gynecologists.

The highest quintiles of maternal weight gain during pregnancy were significantly associated with rates of NICU admission in a study of 2,784 singleton pregnancies, Dr. Teresa Tam and her colleagues, of Saint Joseph Hospital, Chicago, reported in a poster.

After adjusting for age, delivery method, and prepregnancy weight, among other factors, the medium weight gain quintiles—22-29 pounds and 30-35 pounds—were associated with the lowest NICU transfer rates of 3.3% and 2.6%, respectively.

A second poster by Dr. Devendra A. Patel of Weill Cornell Medical College, New York, and colleagues found that heavier women had almost twice the rate of maternal and fetal complications as women of normal weight.

Dr. Patel found no fetal complications and 11 peripartum complications among 68 women whose BMI was less than 30 kg/m<sup>2</sup>, compared with 5 fetal complications and 20 peripartum complications among 78 women whose BMI was 30 kg/m<sup>2</sup> or higher.

—Heidi Splete

**New!**

Can eggs give them added **Fiber?** **Sure!**

**Recommend** clear-mixing, flavor-free **Fibersure**

It's easily added to almost everything patients eat and drink, ensuring fiber compliance.

- Provides 20% of the recommended daily amount of fiber per serving—more than any other leading brand!
- Clear-mixing, non-thickening, flavor-free, 100%-natural fiber
- Mixes, cooks & bakes into most of their favorite foods

Visit [www.fibersure.com](http://www.fibersure.com) or call us at 1-800-525-2855

© 2006 P&G GPAD06011 2/06

from the makers of **Metamucil**

**New!**

**Fiber-sure**

Clear-Mixing Fiber Supplement

Flavor-Free, Grit-Free, Non-Thickening

Try for Baking & Cooking too!

Get them on fiber. Keep them on fiber.